

A SUPERONVERGENCE ANALYSIS FOR FINITE ELEMENT SOLUTION BY THE INTERPOLANT POSTPROCESSING ON IRREGULAR MESHES FOR SMOOTH PROBLEM^{*1)}

Qi-ding Zhu

(Institute of Science, Hunan Normal University, Changsha 410081, China)

Qun Lin

(Institute of System, Academy of Mathematics and Systems Sciences, Chinese Academy of Sciences, Beijing 100080, China)

Abstract

The post-processing procedure is given by a interpolant postprocessing of the finite element solution by appropriately-defined finite dimensional subspaces. The corresponding superconvergence are established on general quasi-regular finite element partitions.

Key words: Finite element, Superconvergence interpolant postprocessing.

1. Introduction

The results in this paper are based on the idea of interpolation postprocessing in [1] and the techniques of L^2 projection processing in [2]

For simplicity, we consider the model problem: Find $u \in H_0^1(\Omega)$, such that

$$\begin{cases} -\nabla \cdot (a\nabla u) = f, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega \end{cases} \quad (1.1)$$

Suppose that J^h and J^H are irregular triangulations (or quadrilateral partitions). Their sizes satisfy $h \ll H$, ($H \rightarrow 0$). Construct piecewise k -order and r -order finite element space S^h and S^H respectively. Let $u^h \in S^h$ be the Galerkin approximation of $u \in H_0^1(\Omega)$, and

$$I_H : C(\bar{\Omega}) \rightarrow S^H \quad (1.2)$$

be the interpolation operator, which satisfies the following three conditions:

- 1) $\|I_H w\|_{1,\infty} \leq CH^{-1}\|I_H w\|_{0,\infty}$,
- 2) $\|u - I_H u\|_{0,\infty} \leq C\|u\|_{0,\infty}, \forall u \in C(\Omega)$
- 3) $\|u - I_H u\|_{m,\infty} \leq CH^{r+1-m}\|u\|_{r+1,\infty}, m = 0, 1$

Obviously the standard Lagrange interpolation operator and the projection interpolation operator proposed in [1] satisfy the above three conditions.

It has been shown in [1] the, if S^h and S^H are 1 or 2 order finite elements of uniform triangulation, or Q^k type elements defined on rectangular partition, then the following nonlocal superconvergence estimation holds when the parameters H and r properly match.

$$\|\nabla(u - I_H u^h)\|_{0,\infty} \leq Ch^{l-\epsilon} \quad (1.3)$$

* Received July 17, 1999.

¹⁾The work supported by the Foundation of National Natural Science Of China and the Foundation of Education of Hunan Province.

where $l = k + 1$ Here we must impose strict assumptions on the partition, so its applicability is limited.

It has been stated that, if we substitute I_H by the L^2 interpolation operator from $L^2(\Omega)$ onto S^H , then for every irregular partitin J^h and J^H we have the following nonlocal L^2 superconvergence estimation, provided that the two parameters H and r properly match:

$$\|\nabla(u - Q_H u^h)\|_0 \leq Ch^{l-\epsilon} [\|u\|_l + \|u\|_{r+1}] \tag{1.4}$$

where $l = k + 1$

The two processing techniques have their own advantages respectivly. The former is very easy to perform interpolation processing, and need not to solve algebraic equations, but strict assumptions must by imposed on the partition. The later is quite contrary.

In this paper, we conclute that ,for every irregular partitions J^H and J^h , if we properly choose the parameters H and r , (1.3) holds. This combines the advantages of the two techniques.

2. Main Results and Proof

Let

$$\bar{k} = \begin{cases} 1, & \text{when } k = 1 \\ 0, & \text{when } k > 1 \end{cases} \tag{2.1}$$

We have the following important Theorem.

Theorem 1. *Suppose that $u \in W^{r+1,\infty}(\Omega) \cap H_0^1(\Omega)$, ($r > k$), $u^h \in S^h$ is k -order Galerkin aprroximation of u , the interpolation operator I_H satisfies the conditions 1),2),3), then we have basic estimation*

$$\|\nabla(u - I_H u^h)\|_{0,\infty} \leq C(H^r \|u\|_{r+1,\infty} + H^{-1} \|u - u^h\|_{0,\infty}) \tag{2.2}$$

or

$$\|\nabla(u - I_H u^h)\|_{0,\infty} \leq C(H^r + H^{-1} h^{k+1} |\log h|^{\bar{k}}) \|u\|_{r+1,\infty} \tag{2.3}$$

Proof. Using the triangular inequality, the condition 3), inverse property 1) and the condition 2) we have

$$\begin{aligned} \|\nabla(u - I_H u^h)\|_{0,\infty} &\leq \|\nabla(u - I_H u)\|_{0,\infty} + \|\nabla(I_H u - I_H u^h)\|_{0,\infty} \\ &\leq CH^r \|u\|_{r+1,\infty} + CH^{-1} \|I_H(u - u^h)\|_{0,\infty} \\ &\leq CH^r \|u\|_{r+1,\infty} + CH^{-1} \|u - u^h\|_{0,\infty} \end{aligned}$$

Then by the well-known L^∞ estimation

$$\|u - u^h\|_{0,\infty} \leq Ch^{k+1} |\log h|^{\bar{k}} \|u\|_{k+1,\infty}$$

we obtain (2.3)

Corollary 1. *Under the conditions of Theorem 1, for every $\epsilon > 0$, there exist positive integer number r and $H \gg h$, such that*

$$\|\nabla(u - I_H u^h)\|_{0,\infty} \leq Ch^{k+1-\epsilon} (\|u\|_{r+1,\infty} + \|u\|_{k+1,\infty}) \tag{2.4}$$

Proof. Select r properly Large so that $\frac{k+1}{r+1} < \epsilon$. Let

$$H = h^{\frac{k+1}{r+1}} \tag{2.5}$$