

PERTURBATION OF ANGLES BETWEEN LINEAR SUBSPACES*

SUN JI-GUANG (孙继广)

(Computing Center, Academia Sinica, Beijing, China)

Abstract

We consider in this note how the principal angles between column spaces $\mathcal{R}(A)$ and $\mathcal{R}(B)$ change when the elements in A and B are subject to perturbations. The basic idea in the proof of our results is that the non-zero cosine values of the principal angles between $\mathcal{R}(A)$ and $\mathcal{R}(B)$ coincide with the non-zero singular values of $P_A P_B$, the product of two orthogonal projections, and consequently we can apply a perturbation theorem of orthogonal projections proved by the author^[4].

§ 1. Introduction

Let \mathcal{X} and \mathcal{Y} be given subspaces of the complex n -dimensional vector space \mathbb{C}^n , and assume that

$$p = \dim(\mathcal{X}) \geq \dim(\mathcal{Y}) = q \geq 1.$$

The principal angles $\theta_k \in [0, \pi/2]$ between \mathcal{X} and \mathcal{Y} are recursively defined for $k=1, \dots, q$ by

$$\cos \theta_k = \max_{u \in \mathcal{X}} \max_{v \in \mathcal{Y}} u^H v = u_k^H v_k, \quad \|u\|_2 = \|v\|_2 = 1,$$

subject to the constraints

$$u_j^H u = 0, \quad v_j^H v = 0, \quad j=1, 2, \dots, k-1.$$

In statistics the numbers $\cos \theta_1, \dots, \cos \theta_k$ are called canonical correlation coefficients. Björck and Golub^[1] pointed out that the principal angles are uniquely defined and have many important applications in statistics and numerical analysis.

Let $\mathcal{R}(A)$ be the column space of a complex $n \times p$ matrix A , and $\mathcal{R}(B)$ be the column space of a complex $n \times q$ matrix B . First order perturbation results for principal angles between $\mathcal{R}(A)$ and $\mathcal{R}(B)$ are given in [1], under the hypotheses of having a small change of A and B . The aim in this note is to remove the hypotheses and to derive perturbation bounds for principal angles. The main results are Theorem 3.1 and Theorem 3.2.

Notation. The symbol $\mathbb{C}^{n \times m}$ denotes the set of complex $n \times m$ matrices, and

$$\mathbb{C}_p^{n \times m} = \{A \in \mathbb{C}^{n \times m} : \text{rank}(A) = p\}.$$

$\sigma(A)$ denotes the set of singular values of a matrix A , and $\sigma_+(A)$ the set of non-zero singular values of A . A^H is for conjugate transpose of A , and $I^{(n)}$ is the $n \times n$ identity matrix. A^\dagger stands for the Moore-Penrose generalized inverse of a matrix A . $P_A = AA^\dagger$ is the orthogonal projection onto the column space $\mathcal{R}(A)$.

* Received October 14, 1985.

Besides, let $\|\cdot\|: \bigcup_{n,m=1}^{\infty} \mathbb{C}^{n \times m} \rightarrow \mathbb{R}_+$ (the set of nonnegative real numbers) be a family of unitarily invariant norms (u. i. n.), $\|\cdot\|_2$ be the spectral norm and $\|\cdot\|_F$ the Frobenius norm.

§ 2. Preliminaries

In this paragraph we will cite some results as the basis of our perturbation theorems for principal angles.

Theorem 2.1^[1]. Let $A \in \mathbb{C}_p^{n \times p}$ and $B \in \mathbb{C}_q^{n \times q}$, $p \geq q$. Assume that the columns of matrices U_A and U_B form unitary bases for two subspaces $\mathcal{R}(A)$ and $\mathcal{R}(B)$. Let the singular value decomposition (SVD) of the $p \times q$ matrix $U_A^H U_B$ be

$$U_A^H U_B = U C V^H, \quad C = \text{diag}(c_1, \dots, c_q),$$

where $U^H U = V^H V = V V^H = I^{(q)}$. If we assume that $c_1 \geq c_2 \geq \dots \geq c_q$, then the principal angles $\theta_1, \dots, \theta_q$ associated with $\mathcal{R}(A)$ and $\mathcal{R}(B)$ are given by

$$\cos \theta_k = c_k, \quad k = 1, \dots, q.$$

In the following we use the symbol $\sigma(A, B)$ for the set $\{c_k\}_{k=1}^q$.

Theorem 2.2^[2]. Let $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_q$ and $\tilde{\sigma}_1 \geq \tilde{\sigma}_2 \geq \dots \geq \tilde{\sigma}_q$ be the singular values of G and $\tilde{G} \in \mathbb{C}^{p \times q}$, respectively, $p \geq q$. Then for every unitarily invariant norm,

$$\|\text{diag}(\tilde{\sigma}_1 - \sigma_1, \dots, \tilde{\sigma}_q - \sigma_q)\| \leq \|\tilde{G} - G\|.$$

Theorem 2.3^[4]. Let $Z, W \in \mathbb{C}^{n \times m}$. If $\text{rank}(W) = \text{rank}(Z)$, then

$$\|P_W - P_Z\| \leq \mu \min\{\|Z^\dagger\|_2, \|W^\dagger\|_2\} \|W - Z\|,$$

where μ is given in the following table:

$\ \cdot\ $	arbitrary u. i. n.	Frobenius	spectral
μ	2	$\sqrt{2}$	1

§ 3. Perturbation Theorems

Let $A \in \mathbb{C}_p^{n \times p}$, $B \in \mathbb{C}_q^{n \times q}$, $p \geq q$. Assume that the columns of U_A and U_B form unitary bases for $\mathcal{R}(A)$ and $\mathcal{R}(B)$, respectively. First we prove two lemmas.

Lemma 3.1. Suppose that $\sigma(U_A^H U_B) = \{c_k\}_{k=1}^q$, $c_k = \cos \theta_k$, $k = 1, \dots, q$, $\frac{\pi}{2} \geq \theta_1 \geq \dots \geq \theta_q \geq 0$. If (U_A, W_A) is an $n \times n$ unitary matrix and $\sigma(W_A^H U_B) = \{s_k\}_{k=1}^q$, $s_1 \geq \dots \geq s_q$, then

$$s_k = \sin \theta_k, \quad k = 1, \dots, q. \quad (3.1)$$

Proof. From

$$(U_A^H U_B)^H (U_A^H U_B) + (W_A^H U_B)^H (W_A^H U_B) = I^{(q)}$$

it follows that

$$c_k^2 + s_k^2 = 1, \quad k = 1, \dots, q.$$

Thus we get the relations (3.1). \blacksquare

Lemma 3.2. For the above mentioned A, B, U_A, U_B and W_A , we have