

Well-Posedness of MHD Equations in Sobolev-Gevery Space

Qian Liu¹ and Baoquan Yuan^{1,†}

Abstract This paper is devoted to the study of the 3D incompressible magnetohydrodynamic system. We prove the local in time well-posedness for any large initial data in $\dot{H}_{a,1}^1(\mathbb{R}^3)$ or $H_{a,1}^1(\mathbb{R}^3)$. Furthermore, the global well-posedness of a strong solution in $\tilde{L}^\infty(0, T; H_{a,1}^1(\mathbb{R}^3)) \cap L^2(0, T; \dot{H}_{a,1}^1(\mathbb{R}^3) \cap \dot{H}_{a,1}^2(\mathbb{R}^3))$ with initial data satisfying a smallness condition is established.

Keywords MHD equation, Sobolev-Gevery space, well-posedness

MSC(2010) 35Q35, 35D35, 76W05.

1. Introduction

The magnetohydrodynamic equations reflect the basic physics laws governing the dynamics of electrically conducting fluids. The velocity field obeys the Navier-Stokes equations, and the magnetic field satisfies the Maxwell's equations of electromagnetism. The magnetohydrodynamic equations play important roles in the study of many phenomena in geophysics, astrophysics, and cosmology(see, [1–3]). In this paper, we consider the 3D incompressible magnetohydrodynamic (short written MHD) equations, which can be written as:

$$\begin{cases} \partial_t u - \mu \Delta u + u \cdot \nabla u + \nabla p = b \cdot \nabla b, \\ \partial_t b - \nu \Delta b + u \cdot \nabla b = b \cdot \nabla u, \\ \nabla \cdot u = 0, \nabla \cdot b = 0, \\ u(x, 0) = u_0(x), b(x, 0) = b_0(x), \end{cases} \quad (1.1)$$

for $t > 0$, $x \in \mathbb{R}^3$. We denote $u = u(x, t)$, $b = b(x, t)$ and $p = p(x, t)$ the velocity field, magnetic field and scalar pressure respectively. The constants μ and ν are the viscosity and resistivity coefficient, u_0 and b_0 are the initial velocity field and initial magnetic field satisfying $\nabla \cdot u_0 = 0$, $\nabla \cdot b_0 = 0$. When $b = 0$, equation (1.1) reduces to the classical Navier-Stokes equation, which has been investigated in many exciting results. Leray [4] and Hopf [5] established the global existence of weak solutions. Fujita and Kato [6] obtained the local well-posedness for large initial data and the global well-posedness for small initial data in Sobolev space. And Kato [7] established similar results in $L^n(\mathbb{R}^n)$. Lei and Lin [8] proved the existence of global mild solution with small initial data in the critical space $\chi^{-1}(\mathbb{R}^3)$. Benameur

[†]the corresponding author.

Email address: liuq971505479@163.com., bqyuan@hpu.edu.cn

¹School of Mathematics and Information Science, Henan Polytechnic University, Henan, 454000, China.

and Jlali [16] studied the long time decay of global solution in Sobolev-Gevery spaces $\dot{H}_{a,\sigma}^1(\mathbb{R}^3)$. Sun and Liu [20] proved that if $u \in C([0, +\infty), \dot{H}_{a,\frac{1}{\alpha}}^{\frac{5}{2}-2\alpha}(\mathbb{R}^3))$ is a global solution of the 3D fractional Navier-Stokes equation, then $\|u(t)\|_{\dot{H}_{a,\frac{1}{\alpha}}^{\frac{5}{2}-2\alpha}(\mathbb{R}^3)}$ decays to zero as time approaches infinity. More results about the solutions to the Navier-Stokes equations in Sobolev-Gevery spaces can be found in ([17–19]). For the MHD equation (1.1), there are several important results. Duvaut and Lions [9] constructed a global weak solution and local strong solution. Sermange and Temam [10] established the local well-posedness of equation (1.1) in Sobolev space for any initial data. Chaabani [21] proved the local in time well-posedness for any large initial data in $H_{a,\sigma}^{\frac{1}{2}}(\mathbb{T}^3)$ as well as global in time well-posedness when initial data satisfies a smallness condition. More researches on the well-posedness for MHD equations can be referred to ([11–14]).

In this paper, we study not only the well-posedness of the local solution in $C([0, T]; \dot{H}_{a,1}^1(\mathbb{R}^3))$ and $C([0, T]; H_{a,1}^1(\mathbb{R}^3))$ for large initial data but also the well-posedness of the global solution in $\tilde{L}^\infty(0, T; H_{a,1}^1(\mathbb{R}^3)) \cap L^2(0, T; \dot{H}_{a,1}^1(\mathbb{R}^3) \cap \dot{H}_{a,1}^2(\mathbb{R}^3))$ for small initial data. We use the Banach contraction mapping principle to prove it. Although it is considered to be valid, the construction of the work space in the process of proof is delicate and not easy. We present it in this paper. Our results are stated in the following theorems.

Theorem 1.1. *Consider the MHD equation (1.1) with $\mu > 0$, and $\nu > 0$. Assume $(u_0, b_0) \in \dot{H}_{a,1}^1(\mathbb{R}^3)$ with $\operatorname{div}u_0 = \operatorname{div}b_0 = 0$. There exists a time $T = T(\|u_0\|_{\dot{H}_{a,1}^1}, \|b_0\|_{\dot{H}_{a,1}^1}) > 0$, such that (1.1) has a unique solution $(u, b) \in C([0, T]; \dot{H}_{a,1}^1(\mathbb{R}^3))$.*

Theorem 1.2. *Consider the MHD equation (1.1) with $\mu > 0$, and $\nu > 0$. Assume $(u_0, b_0) \in H_{a,1}^1(\mathbb{R}^3)$ with $\operatorname{div}u_0 = \operatorname{div}b_0 = 0$. There exists a time $T = T(\|u_0\|_{H_{a,1}^1}, \|b_0\|_{H_{a,1}^1}) > 0$, such that (1.1) has a unique solution $(u, b) \in C([0, T]; H_{a,1}^1(\mathbb{R}^3))$.*

Theorem 1.3. *Consider the MHD equation (1.1) with $\mu > 0$, and $\nu > 0$. Assume $(u_0, b_0) \in H_{a,1}^1(\mathbb{R}^3)$ with $\operatorname{div}u_0 = \operatorname{div}b_0 = 0$. There exists a small enough constant $\varepsilon > 0$ such that if $\|(u_0, b_0)\|_{H_{a,1}^1} \leq \varepsilon$, then system (1.1) has a unique global solution $(u, b) \in \tilde{L}^\infty(0, T; H_{a,1}^1(\mathbb{R}^3)) \cap L^2(0, T; \dot{H}_{a,1}^1(\mathbb{R}^3) \cap \dot{H}_{a,1}^2(\mathbb{R}^3))$, for any $T > 0$.*

2. Notations and lemmas

In this section, we first introduce some notations and definitions that will be used later, then we present several tool lemmas which serve as preparation for the proof of our main results.

- The Fourier transformation is defined as

$$\mathcal{F}(f)(\xi) = \hat{f}(\xi) = \int_{\mathbb{R}^3} e^{-ix \cdot \xi} f(x) dx.$$

- The homogeneous Sobolev space is defined as

$$\dot{H}^s = \{f \in \mathcal{S}'(\mathbb{R}^3); \hat{f} \in L_{loc}^1 \text{ and } |\xi|^s \hat{f} \in L^2(\mathbb{R}^3)\}.$$

- The Sobolev-Gevery space is defined as: for $a > 0$, $s \geq 0$, $\sigma \geq 1$,

$$\dot{H}_{a,\sigma}^s(\mathbb{R}^3) = \{f \in L^2(\mathbb{R}^3) : e^{a\Lambda^{\frac{1}{\sigma}}} f \in \dot{H}^s\},$$

with the norm

$$\|f(t)\|_{\dot{H}_{a,\sigma}^s} = \left(\int_{\mathbb{R}^3} |\xi|^{2s} e^{2a|\xi|^{\frac{1}{\sigma}}} |\hat{f}(t, \xi)|^2 d\xi \right)^{\frac{1}{2}} = \|e^{a\Lambda^{\frac{1}{\sigma}}} f\|_{\dot{H}^s}.$$

Here the Zygmund operator $\Lambda \triangleq (-\Delta)^{\frac{1}{2}}$ is defined via Fourier transform $\widehat{\Lambda f}(\xi) = |\xi| \widehat{f}(\xi)$.

- The Lebesgue-Gevery space $L_{a,\sigma}^p$ is defined as follows. For $1 \leq p \leq \infty$, $a > 0$, $\sigma \geq 1$,

$$L_{a,\sigma}^p(\mathbb{R}^3) = \{f \in L^p(\mathbb{R}^3) : e^{a\Lambda^{\frac{1}{\sigma}}} f \in L^p\},$$

with the norm

$$\|f\|_{L_{a,\sigma}^p} = \left(\int_{\mathbb{R}^3} (e^{a|\xi|^{\frac{1}{\sigma}}} |\hat{f}(t, \xi)|)^p d\xi \right)^{\frac{1}{p}} = \|e^{a\Lambda^{\frac{1}{\sigma}}} f\|_{L^p}.$$

We shall use the Banach contraction mapping principle which can be found in [15], Lemma 5.5.

Lemma 2.1. *Let E be a Banach space, \mathcal{B} a continuous bilinear map from $E \times E$ to E , and α a positive real number such that*

$$\alpha < \frac{1}{4\|\mathcal{B}\|} \text{ with } \|\mathcal{B}\| = \sup_{\|f\|, \|g\| \leq 1} \|\mathcal{B}(f, g)\|.$$

For any a in the ball $B(0, \alpha)$ (with center 0 and radius α) in E , a unique x then exists in $B(0, 2\alpha)$ such that

$$x = a + \mathcal{B}(x, x).$$

The following Lemma 2.2 is a Hölder inequality in the Lebesgue-Gevery spaces.

Lemma 2.2. *Let $a > 0$, $\sigma \geq 1$, $p > 1$, $q > 1$, $\frac{1}{p} + \frac{1}{q} = \frac{1}{2}$. Then*

$$\|u \otimes v\|_{L_{a,\sigma}^2(\mathbb{R}^3)} \leq \|u\|_{L_{a,\sigma}^p(\mathbb{R}^3)} \|v\|_{L_{a,\sigma}^q(\mathbb{R}^3)}. \quad (2.1)$$

Proof.

$$\|u \otimes v\|_{L_{a,\sigma}^2} = \left(\int_{\xi} e^{2a|\xi|^{\frac{1}{\sigma}}} |\widehat{u \otimes v}(\xi)|^2 d\xi \right)^{\frac{1}{2}} \leq \left(\int_{\xi} \left(\int_{\eta} e^{a|\xi|^{\frac{1}{\sigma}}} |\widehat{u}(\xi - \eta)| |\widehat{v}(\eta)| d\eta \right)^2 d\xi \right)^{\frac{1}{2}}. \quad (2.2)$$

Using the inequality $e^{a|\xi|^{\frac{1}{\sigma}}} \leq e^{a|\xi - \eta|^{\frac{1}{\sigma}}} e^{a|\eta|^{\frac{1}{\sigma}}}$, we get

$$\begin{aligned} \|u \otimes v\|_{L_{a,\sigma}^2} &\leq \left(\int_{\xi} \left(\int_{\eta} e^{a|\xi - \eta|^{\frac{1}{\sigma}}} |\widehat{u}(\xi - \eta)| e^{a|\eta|^{\frac{1}{\sigma}}} |\widehat{v}(\eta)| d\eta \right)^2 d\xi \right)^{\frac{1}{2}} \\ &\leq \|UV\|_{L^2} \leq \|U\|_{L^p} \|V\|_{L^q} \leq \|u\|_{L_{a,\sigma}^p} \|v\|_{L_{a,\sigma}^q}, \end{aligned}$$

where $\widehat{U} = e^{a|\xi|^{\frac{1}{\sigma}}} |\widehat{u}(\xi)|$, and $\widehat{V} = e^{a|\xi|^{\frac{1}{\sigma}}} |\widehat{v}(\xi)|$. □

The similar argument can also deduce the following Lemma 2.3.

Lemma 2.3. ([17]) For any $(\alpha, \beta) \in (-\frac{3}{2}, \frac{3}{2})$, a constant C exists such that if $\alpha + \beta$ is positive, then we have

$$\|fg\|_{\dot{H}_{a,1}^{\alpha+\beta-\frac{3}{2}}(\mathbb{R}^3)} \leq C\|f\|_{\dot{H}_{a,1}^{\alpha}(\mathbb{R}^3)}\|g\|_{\dot{H}_{a,1}^{\beta}(\mathbb{R}^3)}. \quad (2.3)$$

The following Lemma 2.4 is the classical $L^p - L^q$ estimates of heat operator.

Lemma 2.4. ([22]) Let $s \geq 0$, $1 \leq p \leq q \leq \infty$. Then

$$\|\Lambda^s e^{t\Delta} f\|_{L^q(\mathbb{R}^d)} \leq C t^{-\frac{s}{2}} t^{-\frac{d}{2}(\frac{1}{p}-\frac{1}{q})} \|f\|_{L^p(\mathbb{R}^d)}. \quad (2.4)$$

The following Lemma 2.5 is an interpolation inequality in the Sobolev-Gevery spaces.

Lemma 2.5. Let $s_1 \leq s \leq s_2$, $a > 0$, $\sigma \geq 1$, $0 < \theta < 1$. Then, $\dot{H}_{a,\sigma}^{s_1} \cap \dot{H}_{a,\sigma}^{s_2}$ is included in $\dot{H}_{a,\sigma}^s$, and we have

$$\|u\|_{\dot{H}_{a,\sigma}^s} \leq \|u\|_{\dot{H}_{a,\sigma}^{s_1}}^{1-\theta} \|u\|_{\dot{H}_{a,\sigma}^{s_2}}^{\theta} \quad \text{with } s = (1-\theta)s_1 + \theta s_2.$$

Proof. Using the Sobolev interpolation inequality which can be found in [15], Proposition 1.32, we have

$$\|f\|_{\dot{H}^s} \leq \|f\|_{\dot{H}^{s_1}}^{1-\theta} \|f\|_{\dot{H}^{s_2}}^{\theta}, \quad s = (1-\theta)s_1 + \theta s_2.$$

Let $f = e^{a\Lambda^{\frac{1}{\sigma}}} u$, and the Lemma 2.5 is proved. \square

The similar argument can also infer the following Lemma 2.6.

Lemma 2.6. Let $a > 0$. Then

$$\|u\|_{L_{a,1}^4} \leq \|u\|_{L_{a,1}^2}^{\frac{1}{4}} \|u\|_{\dot{H}_{a,1}^1}^{\frac{3}{4}}. \quad (2.5)$$

Proof. Using the Lebesgue interpolation inequality and $\|f\|_{L^6} \leq \|f\|_{\dot{H}^1}$, we have

$$\|f\|_{L^4} \leq \|f\|_{L^2}^{\frac{1}{4}} \|f\|_{L^6}^{\frac{3}{4}} \leq \|f\|_{L^2}^{\frac{1}{4}} \|f\|_{\dot{H}^1}^{\frac{3}{4}}.$$

Let $f = e^{a\Lambda} u$, and the Lemma 2.6 is proved. \square

3. Proof of Theorem 1.1

Taking the integral form of system (1.1), we have

$$u(t, x) = e^{t\Delta} u_0(x) - \int_0^t e^{(t-s)\Delta} \mathbb{P}(u \cdot \nabla u)(s) ds + \int_0^t e^{(t-s)\Delta} \mathbb{P}(b \cdot \nabla b)(s) ds, \quad (3.1)$$

$$b(t, x) = e^{t\Delta} b_0(x) - \int_0^t e^{(t-s)\Delta} (u \cdot \nabla b)(s) ds + \int_0^t e^{(t-s)\Delta} (b \cdot \nabla u)(s) ds, \quad (3.2)$$

where $\mathbb{P} := Id - \nabla(-\Delta)^{-1} div$ stands for the Leray projector onto a divergence-free vector field.

We will construct a local solution in $\dot{H}_{a,1}^1(\mathbb{R}^3)$. We first prove the estimate of $\|(u, b)\|_{L^\infty(\dot{H}_{a,1}^1)}$. Applying (3.1) and (3.2) by $\Lambda e^{a\Lambda}$, we get

$$\begin{aligned}\Lambda e^{a\Lambda} u(t, x) &= e^{t\Delta} \Lambda e^{a\Lambda} u_0(x) - \int_0^t \Lambda e^{(t-s)\Delta} e^{a\Lambda} \nabla \mathbb{P}(u \otimes u)(s) ds \\ &\quad + \int_0^t \Lambda e^{(t-s)\Delta} e^{a\Lambda} \nabla \mathbb{P}(b \otimes b)(s) ds \\ &\triangleq I_1 + I_2 + I_3,\end{aligned}\tag{3.3}$$

$$\begin{aligned}\Lambda e^{a\Lambda} b(t, x) &= e^{t\Delta} \Lambda e^{a\Lambda} b_0(x) - \int_0^t \Lambda e^{(t-s)\Delta} \nabla e^{a\Lambda} (u \otimes b)(s) ds \\ &\quad + \int_0^t \Lambda e^{(t-s)\Delta} \nabla e^{a\Lambda} (b \otimes u)(s) ds \\ &\triangleq J_1 + J_2 + J_3.\end{aligned}\tag{3.4}$$

Taking the L^2 -norm, by Lemma 2.4 with $p = q = 2$,

$$\|I_1\|_{L^2} \leq \|u_0\|_{\dot{H}_{a,1}^1},$$

$$\|J_1\|_{L^2} \leq \|b_0\|_{\dot{H}_{a,1}^1}.$$

By Lemmas 2.4 and 2.3 with $\alpha = \beta = 1$, we have

$$\begin{aligned}\|I_2\|_{L^2} &\leq \int_0^t \|\Lambda^{\frac{3}{2}} e^{(t-s)\Delta} \Lambda^{\frac{1}{2}} e^{a\Lambda} \mathbb{P}(u \otimes u)(s)\|_{L^2} ds \\ &\leq C \int_0^t (t-s)^{-\frac{3}{4}} \|\Lambda^{\frac{1}{2}} e^{a\Lambda} \mathbb{P}(u \otimes u)(s)\|_{L^2} ds \\ &\leq C \int_0^t (t-s)^{-\frac{3}{4}} \|u \otimes u\|_{\dot{H}_{a,1}^{\frac{1}{2}}} ds \\ &\leq CT^{\frac{1}{4}} \|u\|_{L^\infty(\dot{H}_{a,1}^1)}^2.\end{aligned}\tag{3.5}$$

By a similar argument it can be deduced

$$\|I_3\|_{L^2} \leq CT^{\frac{1}{4}} \|b\|_{L^\infty(\dot{H}_{a,1}^1)}^2,\tag{3.6}$$

$$\|J_2\|_{L^2} \leq CT^{\frac{1}{4}} \|u\|_{L^\infty(\dot{H}_{a,1}^1)} \|b\|_{L^\infty(\dot{H}_{a,1}^1)},\tag{3.7}$$

and

$$\|J_3\|_{L^2} \leq CT^{\frac{1}{4}} \|b\|_{L^\infty(\dot{H}_{a,1}^1)} \|u\|_{L^\infty(\dot{H}_{a,1}^1)}.\tag{3.8}$$

Thus we obtain

$$\|(u, b)\|_{L^\infty(\dot{H}_{a,1}^1)} \leq \|(u_0, b_0)\|_{\dot{H}_{a,1}^1} + CT^{\frac{1}{4}} \|(u, b)\|_{L^\infty(\dot{H}_{a,1}^1)}^2.\tag{3.9}$$

By Lemma 2.1, take $\|u_0\|_{\dot{H}_{a,1}^1} + \|b_0\|_{\dot{H}_{a,1}^1} \leq \frac{1}{4C_0T^{\frac{1}{4}}}$ for $C_0 > C$. Thus the local existence time $T \leq \left[\frac{1}{4C_0(\|u_0\|_{\dot{H}_{a,1}^1} + \|b_0\|_{\dot{H}_{a,1}^1})} \right]^4$, and we construct a local solution on $[0, T]$.

By (3.1) and (3.2), $u(t, x)$ and $b(t, x)$ are continuous with respect to t , and their $\dot{H}_{a,1}^1(\mathbb{R}^3)$ is bounded. By the Lebesgue dominated convergence theorem, we obtain that $(u, b) \in C\left([0, T]; \dot{H}_{a,1}^1(\mathbb{R}^3)\right)$. The proof of Theorem 1.1 is finished.

4. Proof of Theorem 1.2

We can also construct a local solution in $H_{a,1}^1(\mathbb{R}^3)$. We use the estimate of $\|(u, b)\|_{L^\infty(\dot{H}_{a,1}^1)}$ in section 3. Next, we prove the estimate of $\|(u, b)\|_{L^\infty(L_{a,1}^2)}$. Applying (3.1) and (3.2) by $e^{a\Lambda}$, we get

$$\begin{aligned} e^{a\Lambda}u(t, x) &= e^{t\Delta}e^{a\Lambda}u_0(x) - \int_0^t \nabla e^{(t-s)\Delta}e^{a\Lambda}\mathbb{P}(u \otimes u)(s)ds \\ &\quad + \int_0^t \nabla e^{(t-s)\Delta}e^{a\Lambda}\mathbb{P}(b \otimes b)(s)ds \\ &\triangleq K_1 + K_2 + K_3, \end{aligned} \tag{4.1}$$

$$\begin{aligned} e^{a\Lambda}b(t, x) &= e^{t\Delta}e^{a\Lambda}b_0(x) - \int_0^t \nabla e^{(t-s)\Delta}e^{a\Lambda}(u \otimes b)(s)ds \\ &\quad + \int_0^t \nabla e^{(t-s)\Delta}e^{a\Lambda}(b \otimes u)(s)ds \\ &\triangleq M_1 + M_2 + M_3. \end{aligned} \tag{4.2}$$

Taking the L^2 -norm, we have

$$\begin{aligned} \|K_1\|_{L^2} &\leq \|u_0\|_{L_{a,1}^2}, \\ \|M_1\|_{L^2} &\leq \|b_0\|_{L_{a,1}^2}. \end{aligned}$$

By Lemma 2.4 with $p = q = 2$ and Lemma 2.3 with $\alpha = \beta = 1$, we get

$$\begin{aligned} \|K_2\|_{L^2} &\leq \int_0^t \|\Lambda^{\frac{1}{2}}e^{(t-s)\Delta}\Lambda^{\frac{1}{2}}e^{a\Lambda}\mathbb{P}(u \otimes u)(s)\|_{L^2} ds \\ &\leq C \int_0^t (t-s)^{-\frac{1}{4}} \|\Lambda^{\frac{1}{2}}e^{a\Lambda}\mathbb{P}(u \otimes u)(s)\|_{L^2} ds \\ &\leq C \int_0^t (t-s)^{-\frac{1}{4}} \|u \otimes u\|_{\dot{H}_{a,1}^{\frac{1}{2}}} ds \\ &\leq CT^{\frac{3}{4}} \|u\|_{L^\infty(\dot{H}_{a,1}^1)}^2. \end{aligned} \tag{4.3}$$

Arguing similarly to the above inequality (4.3), we obtain

$$\|K_3\|_{L^2} \leq CT^{\frac{3}{4}} \|b\|_{L^\infty(\dot{H}_{a,1}^1)}^2, \tag{4.4}$$

$$\|M_2\|_{L^2} \leq CT^{\frac{3}{4}} \|u\|_{L^\infty(\dot{H}_{a,1}^1)} \|b\|_{L^\infty(\dot{H}_{a,1}^1)}, \quad (4.5)$$

and

$$\|M_3\|_{L^2} \leq CT^{\frac{3}{4}} \|b\|_{L^\infty(\dot{H}_{a,1}^1)} \|u\|_{L^\infty(\dot{H}_{a,1}^1)}. \quad (4.6)$$

Thus we obtain

$$\|(u, b)\|_{L^\infty(L_{a,1}^2)} \leq \|(u_0, b_0)\|_{L_{a,1}^2} + CT^{\frac{3}{4}} \|(u, b)\|_{L^\infty(\dot{H}_{a,1}^1)}^2. \quad (4.7)$$

Combining the estimates (3.9) and (4.7), we have

$$\|(u, b)\|_{L^\infty(H_{a,1}^1)} \leq \|(u_0, b_0)\|_{H_{a,1}^1} + C(T^{\frac{3}{4}} + T^{\frac{1}{4}}) \|(u, b)\|_{L^\infty(H_{a,1}^1)}^2. \quad (4.8)$$

By Lemma 2.1, take $\|u_0\|_{H_{a,1}^1} + \|b_0\|_{H_{a,1}^1} \leq \frac{1}{4C_0(T^{\frac{3}{4}} + T^{\frac{1}{4}})}$ for $C_0 > C$. Thus we can obtain the local existence time $T(\|u_0\|_{H_{a,1}^1}, \|b_0\|_{H_{a,1}^1})$, and we construct a local solution on $[0, T]$.

By (3.1) and (3.2), $u(t, x)$ and $b(t, x)$ are continuous with respect to t , and their $H_{a,1}^1(\mathbb{R}^3)$ is bounded. By the Lebesgue dominated convergence theorem, we obtain that $(u, b) \in C([0, T]; H_{a,1}^1(\mathbb{R}^3))$. The proof of Theorem 1.2 is finished.

5. Proof of Theorem 1.3

Taking the Fourier transform to the integral form of system (1.1), we have

$$\begin{aligned} \hat{u}(t, \xi) &= e^{-\mu t|\xi|^2} \hat{u}_0 - \int_0^t e^{-\mu(t-s)|\xi|^2} \mathbb{P}i\xi \cdot \widehat{u \otimes u}(s) ds \\ &\quad + \int_0^t e^{-\mu(t-s)|\xi|^2} \mathbb{P}i\xi \cdot \widehat{b \otimes b}(s) ds, \end{aligned} \quad (5.1)$$

$$\begin{aligned} \hat{b}(t, \xi) &= e^{-\nu t|\xi|^2} \hat{b}_0 - \int_0^t e^{-\nu(t-s)|\xi|^2} i\xi \cdot \widehat{u \otimes b}(s) ds \\ &\quad + \int_0^t e^{-\nu(t-s)|\xi|^2} i\xi \cdot \widehat{b \otimes u}(s) ds. \end{aligned} \quad (5.2)$$

We first prove the estimate of $\|(u, b)\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)}$. Multiplying (5.1) and (5.2) by $|\xi|e^{a|\xi|}$, and summing up the resulting equations, we get

$$\begin{aligned} &|\xi|e^{a|\xi|} |\hat{u}(t, \xi)| + |\xi|e^{a|\xi|} |\hat{b}(t, \xi)| \\ &\leq e^{-\mu t|\xi|^2} |\xi|e^{a|\xi|} |\hat{u}_0| + e^{-\nu t|\xi|^2} |\xi|e^{a|\xi|} |\hat{b}_0| + \int_0^t |\xi|e^{-\mu(t-s)|\xi|^2} |\xi|e^{a|\xi|} |\widehat{u \otimes u}(s)| ds \\ &\quad + \int_0^t |\xi|e^{-\mu(t-s)|\xi|^2} |\xi|e^{a|\xi|} |\widehat{b \otimes b}(s)| ds + \int_0^t |\xi|e^{-\nu(t-s)|\xi|^2} |\xi|e^{a|\xi|} |\widehat{u \otimes b}(s)| ds \\ &\quad + \int_0^t |\xi|e^{-\nu(t-s)|\xi|^2} |\xi|e^{a|\xi|} |\widehat{b \otimes u}(s)| ds. \end{aligned} \quad (5.3)$$

Taking the L^∞ -norm in time to (5.3), by Young inequality for convolution, we conclude that

$$\begin{aligned}
& \sup_{0 \leq t < \infty} |\xi| e^{a|\xi|} |\hat{u}(t, \xi)| + \sup_{0 \leq t < \infty} |\xi| e^{a|\xi|} |\hat{b}(t, \xi)| \\
& \leq |\xi| e^{a|\xi|} |\hat{u}_0| + |\xi| e^{a|\xi|} |\hat{b}_0| + \frac{1}{\sqrt{2\mu}} \left(\int_0^t |\xi|^2 e^{2a|\xi|} |\widehat{u \otimes u}(s)|^2 ds \right)^{\frac{1}{2}} \\
& \quad + \frac{1}{\sqrt{2\mu}} \left(\int_0^t |\xi|^2 e^{2a|\xi|} |\widehat{b \otimes b}(s)|^2 ds \right)^{\frac{1}{2}} + \frac{1}{\sqrt{2\nu}} \left(\int_0^t |\xi|^2 e^{2a|\xi|} |\widehat{u \otimes b}(s)|^2 ds \right)^{\frac{1}{2}} \\
& \quad + \frac{1}{\sqrt{2\nu}} \left(\int_0^t |\xi|^2 e^{2a|\xi|} |\widehat{b \otimes u}(s)|^2 ds \right)^{\frac{1}{2}}. \tag{5.4}
\end{aligned}$$

Taking L^2 -norm with respect to ξ for (5.4), we obtain

$$\begin{aligned}
& \|u\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} + \|b\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} \\
& \leq \|u_0\|_{\dot{H}_{a,1}^1} + \|b_0\|_{\dot{H}_{a,1}^1} + \frac{1}{\sqrt{2\mu}} \left(\int_0^t \|u \otimes u\|_{\dot{H}_{a,1}^1}^2(s) ds \right)^{\frac{1}{2}} \\
& \quad + \frac{1}{\sqrt{2\mu}} \left(\int_0^t \|b \otimes b\|_{\dot{H}_{a,1}^1}^2(s) ds \right)^{\frac{1}{2}} + \frac{1}{\sqrt{2\nu}} \left(\int_0^t \|u \otimes b\|_{\dot{H}_{a,1}^1}^2(s) ds \right)^{\frac{1}{2}} \\
& \quad + \frac{1}{\sqrt{2\nu}} \left(\int_0^t \|b \otimes u\|_{\dot{H}_{a,1}^1}^2(s) ds \right)^{\frac{1}{2}}. \tag{5.5}
\end{aligned}$$

By Lemma 2.3 with $\alpha = \beta = \frac{5}{4}$ and Lemma 2.5, we have

$$\begin{aligned}
\left(\int_0^t \|u \otimes u\|_{\dot{H}_{a,1}^1}^2(s) ds \right)^{\frac{1}{2}} & \leq C \left(\int_0^t \|u\|_{\dot{H}_{a,1}^{\frac{5}{4}}}^4(s) ds \right)^{\frac{1}{2}} \\
& \leq C \left(\int_0^t \|u\|_{\dot{H}_{a,1}^1}^2 \|u\|_{\dot{H}_{a,1}^1} \|u\|_{\dot{H}_{a,1}^2}(s) ds \right)^{\frac{1}{2}} \\
& \leq C \|u\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} \left(\int_0^t \|u\|_{\dot{H}_{a,1}^1} \|u\|_{\dot{H}_{a,1}^2}(s) ds \right)^{\frac{1}{2}} \\
& \leq C \|u\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} \|\nabla u\|_{L^2(\dot{H}_{a,1}^1)}. \tag{5.6}
\end{aligned}$$

Using a similar argument, we deduce that

$$\left(\int_0^t \|b \otimes b\|_{\dot{H}_{a,1}^1}^2(s) ds \right)^{\frac{1}{2}} \leq C \|b\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} \|\nabla b\|_{L^2(\dot{H}_{a,1}^1)}, \tag{5.7}$$

$$\begin{aligned}
\left(\int_0^t \|u \otimes b\|_{\dot{H}_{a,1}^1}^2(s) ds \right)^{\frac{1}{2}} & \leq \frac{C}{2} (\|u\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} \|\nabla u\|_{L^2(\dot{H}_{a,1}^1)} \\
& \quad + \|b\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} \|\nabla b\|_{L^2(\dot{H}_{a,1}^1)}), \tag{5.8}
\end{aligned}$$

and

$$\left(\int_0^t \|b \otimes u\|_{\dot{H}_{a,1}^1}^2(s) ds \right)^{\frac{1}{2}} \leq \frac{C}{2} (\|u\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} \|\nabla u\|_{L^2(\dot{H}_{a,1}^1)}$$

$$+ \|b\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} \|\nabla b\|_{L^2(H_{a,1}^1)}. \quad (5.9)$$

Inserting the estimates (5.6)-(5.9) into (5.5), we get

$$\begin{aligned} \|(u, b)\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} &\leq \|(u_0, b_0)\|_{\dot{H}_{a,1}^1} \\ &\quad + \max\left(\frac{C}{\sqrt{2\mu}}, \frac{C}{\sqrt{2\nu}}\right) \|(u, b)\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} \|(\nabla u, \nabla b)\|_{L^2(H_{a,1}^1)}. \end{aligned} \quad (5.10)$$

Next, we prove the estimate of $\|(u, b)\|_{\tilde{L}^\infty(L_{a,1}^2)}$. Multiplying (5.1) and (5.2) by $e^{a|\xi|}$, and summing up the resulting equations, we get

$$\begin{aligned} &e^{a|\xi|} |\hat{u}(t, \xi)| + e^{a|\xi|} |\hat{b}(t, \xi)| \\ &\leq e^{-\mu t|\xi|^2} e^{a|\xi|} |\hat{u}_0| + e^{-\nu t|\xi|^2} e^{a|\xi|} |\hat{b}_0| + \int_0^t |\xi| e^{-\mu(t-s)|\xi|^2} e^{a|\xi|} |\widehat{u \otimes u}(s)| ds \\ &\quad + \int_0^t |\xi| e^{-\mu(t-s)|\xi|^2} e^{a|\xi|} |\widehat{b \otimes b}(s)| ds + \int_0^t |\xi| e^{-\nu(t-s)|\xi|^2} e^{a|\xi|} |\widehat{u \otimes b}(s)| ds \\ &\quad + \int_0^t |\xi| e^{-\nu(t-s)|\xi|^2} e^{a|\xi|} |\widehat{b \otimes u}(s)| ds. \end{aligned} \quad (5.11)$$

Taking the L^∞ -norm in time to (5.11), we conclude that

$$\begin{aligned} &\sup_{0 \leq t < \infty} e^{a|\xi|} |\hat{u}(t, \xi)| + \sup_{0 \leq t < \infty} e^{a|\xi|} |\hat{b}(t, \xi)| \\ &\leq e^{a|\xi|} |\hat{u}_0| + e^{a|\xi|} |\hat{b}_0| + \frac{1}{\sqrt{2\mu}} \left(\int_0^t e^{2a|\xi|} |\widehat{u \otimes u}(s)|^2 ds \right)^{\frac{1}{2}} \\ &\quad + \frac{1}{\sqrt{2\mu}} \left(\int_0^t e^{2a|\xi|} |\widehat{b \otimes b}(s)|^2 ds \right)^{\frac{1}{2}} + \frac{1}{\sqrt{2\nu}} \left(\int_0^t e^{2a|\xi|} |\widehat{u \otimes b}(s)|^2 ds \right)^{\frac{1}{2}} \\ &\quad + \frac{1}{\sqrt{2\nu}} \left(\int_0^t e^{2a|\xi|} |\widehat{b \otimes u}(s)|^2 ds \right)^{\frac{1}{2}}. \end{aligned} \quad (5.12)$$

Taking the L^2 -norm on both sides of (5.12), we have

$$\begin{aligned} &\|u\|_{\tilde{L}^\infty(L_{a,1}^2)} + \|b\|_{\tilde{L}^\infty(L_{a,1}^2)} \\ &\leq \|u_0\|_{L_{a,1}^2} + \|b_0\|_{L_{a,1}^2} + \frac{1}{\sqrt{2\mu}} \left(\int_0^t \|u \otimes u\|_{L_{a,1}^2}^2(s) ds \right)^{\frac{1}{2}} \\ &\quad + \frac{1}{\sqrt{2\mu}} \left(\int_0^t \|b \otimes b\|_{L_{a,1}^2}^2(s) ds \right)^{\frac{1}{2}} + \frac{1}{\sqrt{2\nu}} \left(\int_0^t \|u \otimes b\|_{L_{a,1}^2}^2(s) ds \right)^{\frac{1}{2}} \\ &\quad + \frac{1}{\sqrt{2\nu}} \left(\int_0^t \|b \otimes u\|_{L_{a,1}^2}^2(s) ds \right)^{\frac{1}{2}}. \end{aligned} \quad (5.13)$$

Using Lemma 2.2 with $p = q = 4$ and Lemma 2.6, we have

$$\left(\int_0^t \|u \otimes u\|_{L_{a,1}^2}^2(s) ds \right)^{\frac{1}{2}} \leq C \left(\int_0^t \|u\|_{L_{a,1}^4}^4(s) ds \right)^{\frac{1}{2}}$$

$$\begin{aligned}
&\leq C \left(\int_0^t \left(\|u\|_{\dot{H}_{a,1}^1}^{\frac{3}{4}} \|u\|_{L_{a,1}^2}^{\frac{1}{4}} \right)^4 (s) \, ds \right)^{\frac{1}{2}} \\
&\leq C \|u\|_{\tilde{L}^\infty(H_{a,1}^1)} \left(\int_0^t \|u\|_{\dot{H}_{a,1}^1}^2 (s) \, ds \right)^{\frac{1}{2}} \\
&\leq C \|u\|_{\tilde{L}^\infty(H_{a,1}^1)} \|\nabla u\|_{L^2(L_{a,1}^2)}. \tag{5.14}
\end{aligned}$$

By an argument similar to (5.14), we obtain

$$\left(\int_0^t \|b \otimes b\|_{L_{a,1}^2}^2 (s) \, ds \right)^{\frac{1}{2}} \leq C \|b\|_{\tilde{L}^\infty(H_{a,1}^1)} \|\nabla b\|_{L^2(L_{a,1}^2)}, \tag{5.15}$$

$$\begin{aligned}
\left(\int_0^t \|u \otimes b\|_{L_{a,1}^2}^2 (s) \, ds \right)^{\frac{1}{2}} &\leq \frac{C}{2} (\|u\|_{\tilde{L}^\infty(H_{a,1}^1)} \|\nabla u\|_{L^2(L_{a,1}^2)} \\
&\quad + \|b\|_{\tilde{L}^\infty(H_{a,1}^1)} \|\nabla b\|_{L^2(L_{a,1}^2)}), \tag{5.16}
\end{aligned}$$

and

$$\begin{aligned}
\left(\int_0^t \|b \otimes u\|_{L_{a,1}^2}^2 (s) \, ds \right)^{\frac{1}{2}} &\leq \frac{C}{2} (\|u\|_{\tilde{L}^\infty(H_{a,1}^1)} \|\nabla u\|_{L^2(L_{a,1}^2)} \\
&\quad + \|b\|_{\tilde{L}^\infty(H_{a,1}^1)} \|\nabla b\|_{L^2(L_{a,1}^2)}). \tag{5.17}
\end{aligned}$$

Inserting the estimates (5.14)-(5.17) into (5.13), we get

$$\begin{aligned}
\|(u, b)\|_{\tilde{L}^\infty(L_{a,1}^2)} &\leq \|(u_0, b_0)\|_{L_{a,1}^2} \\
&\quad + \max\left(\frac{C}{\sqrt{2\mu}}, \frac{C}{\sqrt{2\nu}}\right) \|(u, b)\|_{\tilde{L}^\infty(H_{a,1}^1)} \|(\nabla u, \nabla b)\|_{L^2(L_{a,1}^2)}. \tag{5.18}
\end{aligned}$$

Combining the estimates (5.10) and (5.18), we have

$$\begin{aligned}
\|(u, b)\|_{\tilde{L}^\infty(H_{a,1}^1)} &\leq \|(u_0, b_0)\|_{H_{a,1}^1} \\
&\quad + \max\left(\frac{C}{\sqrt{2\mu}}, \frac{C}{\sqrt{2\nu}}\right) \|(u, b)\|_{\tilde{L}^\infty(H_{a,1}^1)} \|(\nabla u, \nabla b)\|_{L^2(H_{a,1}^1)}. \tag{5.19}
\end{aligned}$$

To close the estimate, we prove the estimate of $\|(\nabla u, \nabla b)\|_{L^2(0,T;H_{a,1}^1)}$. Firstly, we estimate $\|(\nabla u, \nabla b)\|_{L^2(0,T;\dot{H}_{a,1}^1)}$ which is equal to $\|(u, b)\|_{L^2(0,T;\dot{H}_{a,1}^2)}$. Multiplying (5.1) and (5.2) by $|\xi|^2 e^{a|\xi|}$, we have

$$\begin{aligned}
&|\xi|^2 e^{a|\xi|} |\hat{u}(t, \xi)| + |\xi|^2 e^{a|\xi|} |\hat{b}(t, \xi)| \\
&\leq e^{-\mu t |\xi|^2} |\xi|^2 e^{a|\xi|} |\hat{u}_0| + e^{-\nu t |\xi|^2} |\xi|^2 e^{a|\xi|} |\hat{b}_0| + \int_0^t |\xi|^2 e^{-\mu(t-s)|\xi|^2} |\xi| e^{a|\xi|} |\widehat{u \otimes u}(s)| \, ds \\
&\quad + \int_0^t |\xi|^2 e^{-\mu(t-s)|\xi|^2} |\xi| e^{a|\xi|} |\widehat{b \otimes b}(s)| \, ds + \int_0^t |\xi|^2 e^{-\nu(t-s)|\xi|^2} |\xi| e^{a|\xi|} |\widehat{u \otimes b}(s)| \, ds \\
&\quad + \int_0^t |\xi|^2 e^{-\nu(t-s)|\xi|^2} |\xi| e^{a|\xi|} |\widehat{b \otimes u}(s)| \, ds. \tag{5.20}
\end{aligned}$$

Taking the L^2 -norm with respect to time t in (5.20), we deduce that

$$\begin{aligned}
& \left(\int_0^t |\xi|^4 e^{2a|\xi|} |\hat{u}(t, \xi)|^2 ds \right)^{\frac{1}{2}} + \left(\int_0^t |\xi|^4 e^{2a|\xi|} |\hat{b}(t, \xi)|^2 ds \right)^{\frac{1}{2}} \\
& \leq \frac{1}{\sqrt{2\mu}} |\xi| e^{a|\xi|} |\hat{u}_0| + \frac{1}{\sqrt{2\nu}} |\xi| e^{a|\xi|} |\hat{b}_0| + \frac{1}{\mu} \left(\int_0^t |\xi|^2 e^{2a|\xi|} |\widehat{u \otimes u}(s)|^2 ds \right)^{\frac{1}{2}} \\
& \quad + \frac{1}{\mu} \left(\int_0^t |\xi|^2 e^{2a|\xi|} |\widehat{b \otimes b}(s)|^2 ds \right)^{\frac{1}{2}} + \frac{1}{\nu} \left(\int_0^t |\xi|^2 e^{2a|\xi|} |\widehat{u \otimes b}(s)|^2 ds \right)^{\frac{1}{2}} \\
& \quad + \frac{1}{\nu} \left(\int_0^t |\xi|^2 e^{2a|\xi|} |\widehat{b \otimes u}(s)|^2 ds \right)^{\frac{1}{2}}. \tag{5.21}
\end{aligned}$$

Taking the L^2 -norm with respect to ξ on both sides of (5.21), by an argument similar to (5.10), we have

$$\begin{aligned}
\|(\nabla u, \nabla b)\|_{L^2(\dot{H}_{a,1}^1)} & \leq \max\left(\frac{1}{\sqrt{2\mu}}, \frac{1}{\sqrt{2\nu}}\right) \|(u_0, b_0)\|_{\dot{H}_{a,1}^1} \\
& \quad + \max\left(\frac{C}{\mu}, \frac{C}{\nu}\right) \|(u, b)\|_{\tilde{L}^\infty(\dot{H}_{a,1}^1)} \|(\nabla u, \nabla b)\|_{L^2(H_{a,1}^1)}. \tag{5.22}
\end{aligned}$$

In the following, we estimate the norm $\|(\nabla u, \nabla b)\|_{L^2(0,T;L_{a,1}^2)}$ which is equal to the norm $\|(u, b)\|_{L^2(0,T;\dot{H}_{a,1}^1)}$. Multiplying (5.1) and (5.2) by $|\xi| e^{a|\xi|}$, and summing up the resulting equations, we get

$$\begin{aligned}
& |\xi| e^{a|\xi|} |\hat{u}(t, \xi)| + |\xi| e^{a|\xi|} |\hat{b}(t, \xi)| \\
& \leq e^{-\mu t} |\xi|^2 |\xi| e^{a|\xi|} |\hat{u}_0| + e^{-\nu t} |\xi|^2 |\xi| e^{a|\xi|} |\hat{b}_0| + \int_0^t |\xi|^2 e^{-\mu(t-s)} |\xi|^2 e^{a|\xi|} |\widehat{u \otimes u}(s)| ds \\
& \quad + \int_0^t |\xi|^2 e^{-\mu(t-s)} |\xi|^2 e^{a|\xi|} |\widehat{b \otimes b}(s)| ds + \int_0^t |\xi|^2 e^{-\nu(t-s)} |\xi|^2 e^{a|\xi|} |\widehat{u \otimes b}(s)| ds \\
& \quad + \int_0^t |\xi|^2 e^{-\nu(t-s)} |\xi|^2 e^{a|\xi|} |\widehat{b \otimes u}(s)| ds. \tag{5.23}
\end{aligned}$$

Taking the L^2 -norm with respect to time t in (5.23), we deduce that

$$\begin{aligned}
& \left(\int_0^t |\xi|^2 e^{2a|\xi|} |\hat{u}(t, \xi)|^2 ds \right)^{\frac{1}{2}} + \left(\int_0^t |\xi|^2 e^{2a|\xi|} |\hat{b}(t, \xi)|^2 ds \right)^{\frac{1}{2}} \\
& \leq \frac{1}{\sqrt{2\mu}} e^{a|\xi|} |\hat{u}_0| + \frac{1}{\sqrt{2\nu}} e^{a|\xi|} |\hat{b}_0| + \frac{1}{\mu} \left(\int_0^t e^{2a|\xi|} |\widehat{u \otimes u}(s)|^2 ds \right)^{\frac{1}{2}} \\
& \quad + \frac{1}{\mu} \left(\int_0^t e^{2a|\xi|} |\widehat{b \otimes b}(s)|^2 ds \right)^{\frac{1}{2}} + \frac{1}{\nu} \left(\int_0^t e^{2a|\xi|} |\widehat{u \otimes b}(s)|^2 ds \right)^{\frac{1}{2}} \\
& \quad + \frac{1}{\nu} \left(\int_0^t e^{2a|\xi|} |\widehat{b \otimes u}(s)|^2 ds \right)^{\frac{1}{2}}. \tag{5.24}
\end{aligned}$$

Taking the L^2 -norm on both sides of (5.24), arguing similarly to (5.18), we have

$$\|(\nabla u, \nabla b)\|_{L^2(L_{a,1}^2)} \leq \max\left(\frac{1}{\sqrt{2\mu}}, \frac{1}{\sqrt{2\nu}}\right) \|(u_0, b_0)\|_{L_{a,1}^2}$$

$$+ \max\left(\frac{C}{\mu}, \frac{C}{\nu}\right) \|(u, b)\|_{\tilde{L}^\infty(H_{a,1}^1)} \|(\nabla u, \nabla b)\|_{L^2(L_{a,1}^2)}. \quad (5.25)$$

Combining the estimates (5.22) and (5.25), it arrives at

$$\begin{aligned} \|(\nabla u, \nabla b)\|_{L^2(H_{a,1}^1)} &\leq \max\left(\frac{1}{\sqrt{2\mu}}, \frac{1}{\sqrt{2\nu}}\right) \|(u_0, b_0)\|_{H_{a,1}^1} \\ &+ \max\left(\frac{C}{\mu}, \frac{C}{\nu}\right) \|(u, b)\|_{\tilde{L}^\infty(H_{a,1}^1)} \|(\nabla u, \nabla b)\|_{L^2(H_{a,1}^1)}. \end{aligned} \quad (5.26)$$

Combining the estimates (5.19) and (5.26), we have

$$\begin{aligned} &\|(u, b)\|_{\tilde{L}^\infty(H_{a,1}^1)} + \|(\nabla u, \nabla b)\|_{L^2(H_{a,1}^1)} \\ &\leq \max\left(1, \frac{1}{\sqrt{2\mu}}, \frac{1}{\sqrt{2\nu}}\right) \|(u_0, b_0)\|_{H_{a,1}^1} \\ &+ C \max\left(\frac{1}{\sqrt{2\mu}}, \frac{1}{\mu}, \frac{1}{\sqrt{2\nu}}, \frac{1}{\nu}\right) \|(u, b)\|_{\tilde{L}^\infty(H_{a,1}^1)} \|(\nabla u, \nabla b)\|_{L^2(H_{a,1}^1)}. \end{aligned} \quad (5.27)$$

Applying Lemma 2.1, if we take the initial data small enough,

$$\|(u_0, b_0)\|_{H_{a,1}^1} \leq \frac{1}{4C_0 \max\left(1, \frac{1}{\sqrt{2\mu}}, \frac{1}{\sqrt{2\nu}}\right) \max\left(\frac{1}{\sqrt{2\mu}}, \frac{1}{\mu}, \frac{1}{\sqrt{2\nu}}, \frac{1}{\nu}\right)} = \varepsilon,$$

for $C_0 > C$, then we have a unique global solution $(u, b) \in \tilde{L}^\infty(0, T; H_{a,1}^1(\mathbb{R}^3)) \cap L^2\left(0, T; \dot{H}_{a,1}^1(\mathbb{R}^3) \cap \dot{H}_{a,1}^2(\mathbb{R}^3)\right)$, thus we complete the proof of Theorem 1.3.

References

- [1] D. Biskamp, *Nonlinear Magnetohydrodynamics*, Cambridge University Press, Cambridge, 1993.
- [2] P. A. Davidson, *An introduction to Magnetohydrodynamics*, Cambridge University Press, Cambridge, England, 2001.
- [3] E. Priest and T. Forbes, *Magnetic Reconnection, MHD Theory and Applications*, Cambridge University Press, Cambridge, 2000.
- [4] J. Leray, *Sur le mouvement d'un liquide visqueux emplissant l'espace*, Acta mathematica, 1934, 63, 193-248.
- [5] E. Hopf, *Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen*, Mathematische Nachrichten, 1951, 4, 213-231.
- [6] H. Fujita and T. Kato, *On the Navier-Stokes initial value problem*, Archive for Rational Mechanics and Analysis, 1964, 16(4), 269-315.
- [7] T. Kato, *Strong L^p -solutions of the Navier-Stokes initial in \mathbb{R}^m with applications to weak solutions*, Mathematische Zeitschrift, 1984, 187, 471-480.
- [8] Z. Lei and F. H. Lin, *Global Mild Solutions of the Navier-Stokes Equations*, Communications on Pure and Applied Mathematics, 2011, 64, 1297-1304.
- [9] G. Duvaut and J.-L. Lions, *Inéquations en thermoélasticité et magnétohydrodynamique*, Archive for Rational Mechanics and Analysis, 1972, 46, 241-279.

-
- [10] M. Sermange and R. Temam, *Some mathematical questions related to the MHD equations*, Communications on Pure and Applied Mathematics, 1983, 36, 635-664.
- [11] C. X. Miao, B. Q. Yuan and B. Zhang, *Well-posedness for the incompressible magnetohydrodynamic system*, Mathematical Methods in the Applied Sciences, 2010,30(8), 961-976.
- [12] H. Kozono, *On Weak and Classical Solutions of the Two-Dimensional Magnetohydrodynamic Equations*, Tohoku Mathematical Journal, 1989, 44(1), 83-86.
- [13] Y. Z. Wang and K. Y. Wang, *Global well-posedness of the three dimensional magnetohydrodynamics equations*, Nonlinear Analysis Real World Applications, 2014, 17 245-251.
- [14] Y. M. Xiao and B. Q. Yuan, *Temporal decay of a global solution to 3D magnetohydrodynamic system in critical spaces*, Jqournal Evolution Equations, 2021, 21, 1-15.
- [15] H. Bahouri, J. -Y. Chemin and R. Danchin, *Fourier analysis and nonlinear partial differential equations*, Grundlehren der Mathematischen Wissenschaften, Springer-Verlag, Heidelberg, 2011.
- [16] J. Benameur and L. Jlali, *Long time decay for 3D Navier-Stokes equations in Sobolev-Gevrey spaces*, Electronic Journal of Differential Equations, 2016, 104, 1-13.
- [17] J. Benameur and L. Jlali, *On the blow-up criterion of 3D-NSE in Sobolev-Gevrey spaces*, Journal of Mathematical Fluid Mechanics, 2016, 18(4), 805-822.
- [18] H. Orf, *Long time decay for global solutions to the Navier-Stokes equations in Sobolev-Gevrey spaces*, arXiv: 1903. 03034v1, 2019.
- [19] P. Braz e Silva, W. G. Melo and N. F. Rocha, *Existence, uniqueness and blow-up of solutions for the 3D Navier-Stokes equations in homogeneous Sobolev-Gevrey spaces*, Computational and Applied Mathematics, 2020, 39-66.
- [20] X. C. Sun and J. Liu, *Long time decay of the fractional Navier-Stokes equations in Sobolev-Gevrey spaces*, Journal of Nonlinear Evolution Equations and Applications, 2021, 6, 119-135.
- [21] C. Chaabani, *On well-posedness of the Cauchy problem for 3D MHD system in critical Sobolev-Gevrey space*, SN Partial Differential Equations and Applications, 2021, 2-24.
- [22] R. H. Ji, J. H. Wu and W. R. Yang, *Stability and optimal decay for the 3D Navier-Stokes equations with horizontal dissipation*, Journal of Differential Equations, 2021, 290, 57-77.