

## Nonlinear Degenerate Anisotropic Elliptic Equations with Variable Exponents and $L^1$ data

KHELIFI Hichem<sup>1,2,\*</sup> and MOKHTARI Fares<sup>1</sup>

<sup>1</sup>Department of Mathematics and Informatics, University of Algiers, Algiers, Algeria. 2 Street Didouche Mourad Algiers.

<sup>2</sup> Applied Mathematics Laboratory, Badji Mokhtar University-Annaba B.P. 12, Algeria.

Received 31 March 2019; Accepted 24 January 2020

---

**Abstract.** This paper is devoted to the study of a nonlinear anisotropic elliptic equation with degenerate coercivity, lower order term and  $L^1$  datum in appropriate anisotropic variable exponents Sobolev spaces. We obtain the existence of distributional solutions.

**AMS Subject Classifications:** 35J70, 35J60, 35D30

**Chinese Library Classifications:** O175.27

**Key Words:** Sobolev spaces with variable exponents; anisotropic equations; elliptic equations;  $L^1$  data.

---

### 1 Introduction

In this paper we prove the existence of solutions to the nonlinear anisotropic degenerate elliptic equations with variable exponents, of the type

$$\begin{aligned} -\sum_{i=1}^N D_i a_i(x, u, \nabla u) + g(x, u, \nabla u) &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \partial\Omega, \end{aligned} \tag{1.1}$$

where  $\Omega \subseteq \mathbb{R}^N$  ( $N \geq 3$ ) is a bounded domain with smooth boundary  $\partial\Omega$  and the right-hand side  $f$  in  $L^1(\Omega)$ ,  $D_i u = \frac{\partial u}{\partial x_i}$ . We suppose that  $a_i : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$ ,  $i = 1, \dots, N$  are

---

\*Corresponding author. Email addresses: khelifi.hichemedp@gmail.com (H. Khelifi), f.mokhtari@univ-alger.dz (F. Mokhtari)

Carathéodory functions such that for almost every  $x$  in  $\Omega$  and for every  $(\sigma, \xi) \in \mathbb{R} \times \mathbb{R}^N$  the following assumptions are satisfied for all  $i = 1, \dots, N$

$$|a_i(x, \sigma, \xi)| \leq \beta \left( |k(x)| + |\sigma|^{\bar{p}(x)} + \sum_{j=1}^N |\xi_j|^{p_j(x)} \right)^{1 - \frac{1}{p_i(x)}}, \quad (1.2)$$

$$\sum_{i=1}^N (a_i(x, \sigma, \xi) - a_i(x, \sigma, \eta)) (\xi_i - \eta_i) > 0, \quad \forall \xi \neq \eta, \quad (1.3)$$

$$\sum_{i=1}^N a_i(x, \sigma, \xi) \xi_i \geq \alpha \sum_{i=1}^N \frac{|\xi_i|^{p_i(x)}}{(1 + |\sigma|)^{\gamma_i(x)}}, \quad (1.4)$$

where  $\beta > 0$ ,  $\alpha > 0$ , and  $k \in L^1(\Omega)$ ,  $\gamma_i: \bar{\Omega} \rightarrow \mathbb{R}^+$ ,  $p_i: \bar{\Omega} \rightarrow (1, +\infty)$  are continuous functions and  $\bar{p}$  is such that

$$\frac{1}{\bar{p}(\cdot)} = \frac{1}{N} \sum_{i=1}^N \frac{1}{p_i(\cdot)}.$$

We introduce the function

$$\bar{p}^*(x) = \begin{cases} \frac{N\bar{p}(x)}{N-\bar{p}(x)}, & \text{if } \bar{p}(x) < N \\ +\infty, & \text{if } \bar{p}(x) \geq N. \end{cases} \quad (1.5)$$

The nonlinear term  $g: \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$  is a Carathéodory function such that for a.e.  $x \in \Omega$  and all  $(\sigma, \xi) \in \mathbb{R} \times \mathbb{R}^N$ , we have

$$|g(x, \sigma, \xi)| \leq b(|\sigma|) \left( c(x) + \sum_{i=1}^N |\xi_i|^{p_i(x)} \right), \quad (1.6)$$

$$g(x, \sigma, \xi) \cdot \sigma \geq 0, \quad (1.7)$$

where  $b: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a continuous and increasing function with finite values,  $c \in L^1(\Omega)$  and  $\exists \rho > 0$  such that:

$$|g(x, \sigma, \xi)| \geq \rho \left( \sum_{i=1}^N |\xi_i|^{p_i(x)} \right), \quad \forall \sigma \text{ such that } |\sigma| > \rho. \quad (1.8)$$

In [1], the authors obtain the existence of renormalized and entropy solutions for the nonlinear elliptic equation with degenerate coercivity of the type

$$-\operatorname{div}[a(x, u)|\nabla u|^{p(x)-2}\nabla u] + g(x, u) = f \in L^1(\Omega).$$

For  $g \equiv 0$  and  $f \in L^{m(\cdot)}(\Omega)$ , with  $m(x) \geq m_- \geq 1$ , equation of the form (1.1) have been widely studied in [2], where the authors obtain some existence and regularity results for the solutions. If  $g \equiv |u|^{s(x)-1}u$ ,

$$a_i(x, u, \nabla u) = \frac{|D_i u|^{p_i(x)-2} D_i u}{(1 + |u|)^{\gamma_i(x)}}$$