

1

2 Multiple Integral Inequalities for Schur Convex 3 Functions on Symmetric and Convex Bodies

4 Silvestru Sever Dragomir^{1,2,*}

5 ¹ Mathematics, College of Engineering & Science, Victoria University, Melbourne
6 City, MC 8001, Australia

7 ² DST-NRF Centre of Excellence in the Mathematical, and Statistical Sciences, School
8 of Computer Science, & Applied Mathematics, University of the Witwatersrand,
9 Private Bag 3, Johannesburg 2050, South Africa

10 Received 18 July 2019; Accepted (in revised version) 31 January 2021

12

Abstract. In this paper, by making use of Divergence theorem for multiple integrals, we establish some integral inequalities for Schur convex functions defined on bodies $B \subset \mathbb{R}^n$ that are symmetric, convex and have nonempty interiors. Examples for three dimensional balls are also provided.

13 **Key Words:** Schur convex functions, Multiple integral inequalities.

14 **AMS Subject Classifications:** 26D15

15

16 1 Introduction

For any $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, let $x_{[1]} \geq \dots \geq x_{[n]}$ denote the components of x in decreasing order, and let $x_{\downarrow} = (x_{[1]}, \dots, x_{[n]})$ denote the decreasing rearrangement of x . For $x, y \in \mathbb{R}^n$, $x \prec y$ if, by definition,

$$\begin{cases} \sum_{i=1}^k x_{[i]} \leq \sum_{i=1}^k y_{[i]}, & k = 1, \dots, n-1, \\ \sum_{i=1}^n x_{[i]} = \sum_{i=1}^n y_{[i]}. \end{cases}$$

17 When $x \prec y$, x is said to be majorized by y (y majorizes x). This notation and terminology
18 was introduced by Hardy, Littlewood and Pólya in 1934.

19 Functions that preserve the ordering of majorization are said to be Schur-convex. Per-
20 haps Schur-increasing would be more appropriate, but the term Schur-convex is by now
21 well entrenched in the literature, [5, p. 80].

*Corresponding author. Email address: sever.dragomir@vu.edu.au (S. Dragomir)

22 A real-valued function ϕ defined on a set $\mathcal{A} \subset \mathbb{R}^n$ is said to be Schur-convex on \mathcal{A} if

$$x \prec y \quad \text{on } \mathcal{A} \Rightarrow \phi(x) \leq \phi(y). \quad (1.1)$$

23 If, in addition, $\phi(x) < \phi(y)$ whenever $x \prec y$ but x is not a permutation of y , then ϕ is
 24 said to be strictly Schur-convex on \mathcal{A} . If $\mathcal{A} = \mathbb{R}^n$, then ϕ is simply said to be Schur-convex
 25 or strictly Schur-convex.

26 For fundamental properties of Schur convexity see the monograph [5] and the refer-
 27 ences therein. For some recent results, see [2–4] and [6–8].

28 The following result is known in the literature as Schur-Ostrowski theorem [5, p. 84]:

29 **Theorem 1.1.** *Let $I \subset \mathbb{R}$ be an open interval and let $\phi : I^n \rightarrow \mathbb{R}$ be continuously differentiable.*
 30 *Necessary and sufficient conditions for ϕ to be Schur-convex on I^n are*

$$\phi \text{ is symmetric on } I^n, \quad (1.2)$$

31 and for all $i \neq j$, with $i, j \in \{1, \dots, n\}$,

$$(z_i - z_j) \left[\frac{\partial \phi(z)}{\partial x_i} - \frac{\partial \phi(z)}{\partial x_j} \right] \geq 0 \quad \text{for all } z \in I^n, \quad (1.3)$$

32 where $\frac{\partial \phi}{\partial x_k}$ denotes the partial derivative of ϕ with respect to its k -th argument.

33 With the aid of (1.2), condition (1.3) can be replaced by the condition

$$(z_1 - z_2) \left[\frac{\partial \phi(z)}{\partial x_1} - \frac{\partial \phi(z)}{\partial x_2} \right] \geq 0 \quad \text{for all } z \in I^n. \quad (1.4)$$

34 This simplified condition is sometimes more convenient to verify.

35 The above condition is not sufficiently general for all applications because the domain
 36 of ϕ may not be a Cartesian product.

37 Let $\mathcal{A} \subset \mathbb{R}^n$ be a set with the following properties:

- 38 (i) \mathcal{A} is symmetric in the sense that $x \in \mathcal{A} \Rightarrow x\Pi \in \mathcal{A}$ for all permutations Π ;
 39 (ii) \mathcal{A} is convex and has a nonempty interior.

40 We have the following result, [5, p. 85].

41 **Theorem 1.2.** *If ϕ is continuously differentiable on the interior of \mathcal{A} and continuous on \mathcal{A} , then*
 42 *necessary and sufficient conditions for ϕ to be Schur-convex on \mathcal{A} are*

$$\phi \text{ is symmetric on } \mathcal{A} \quad (1.5)$$

43 and

$$(z_1 - z_2) \left[\frac{\partial \phi(z)}{\partial x_1} - \frac{\partial \phi(z)}{\partial x_2} \right] \geq 0 \quad \text{for all } z \in \mathcal{A}. \quad (1.6)$$

It is well known that any symmetric convex function defined on a symmetric convex set \mathcal{A} is Schur convex, [5, p. 97]. If the function $\phi : \mathcal{A} \rightarrow \mathbb{R}$ is symmetric and quasi-convex, namely

$$\phi(\alpha u + (1 - \alpha)v) \leq \max\{\phi(u), \phi(v)\}$$

44 for all $\alpha \in [0, 1]$ and $u, v \in \mathcal{A}$, a symmetric convex set, then ϕ is Schur convex on \mathcal{A} [5,
45 p. 98].

46 In the recent paper [3] we obtained the following result for Schur convex functions
47 defined on symmetric convex domains of \mathbb{R}^2 .

48 **Theorem 1.3.** *Let $D \subset \mathbb{R}^2$ be symmetric, convex and has a nonempty interior. If ϕ is continu-*
49 *ously differentiable on the interior of D , continuous and Schur convex on D and ∂D is a simple,*
50 *closed counterclockwise curve in the xy -plane bounding D , then*

$$\iint_D \phi(x, y) \, dx dy \leq \frac{1}{2} \oint_{\partial D} [(x - y) \phi(x, y) \, dx + (x + y) \phi(x, y) \, dy]. \quad (1.7)$$

51 *If ϕ is Schur concave on D , then the sign of inequality reverses in (1.7).*

52 Motivated by the above results, we establish in this paper a generalization of the
53 inequality (1.7) for the case of symmetric and convex subsets in n -dimensional space \mathbb{R}^n .
54 This is done by employing an identity obtained via the well known Divergence Theorem
55 for volume and surface integrals. An example for balls in three dimensional space are
56 also provided.

57 2 Some preliminary facts

58 Let B be a bounded open subset of \mathbb{R}^n ($n \geq 2$) with smooth (or piecewise smooth) bound-
59 ary ∂B . Let $F = (F_1, \dots, F_n)$ be a smooth vector field defined in \mathbb{R}^n , or at least in $B \cup \partial B$.
60 Let \mathbf{n} be the unit outward-pointing normal of ∂B . Then the Divergence Theorem states,
61 see for instance [9]:

$$\int_B \operatorname{div} F \, dV = \int_{\partial B} F \cdot \mathbf{n} \, dA, \quad (2.1)$$

where

$$\operatorname{div} F = \nabla \cdot F = \sum_{k=1}^n \frac{\partial F_k}{\partial x_k},$$

62 dV is the element of volume in \mathbb{R}^n and dA is the element of surface area on ∂B .

63 If $\mathbf{n} = (\mathbf{n}_1, \dots, \mathbf{n}_n)$, $x = (x_1, \dots, x_n) \in B$ and use the notation dx for dV we can write
64 (2.1) more explicitly as

$$\sum_{k=1}^n \int_B \frac{\partial F_k(x)}{\partial x_k} \, dx = \sum_{k=1}^n \int_{\partial B} F_k(x) \mathbf{n}_k(x) \, dA. \quad (2.2)$$

65 By taking the real and imaginary part, we can extend the above equality for complex
66 valued functions $F_k, k \in \{1, \dots, n\}$ defined on B .

If $n = 2$, the normal is obtained by rotating the tangent vector through 90° (in the correct direction so that it points out). The quantity $t ds$ can be written (dx_1, dx_2) along the surface, so that

$$\mathbf{n} dA := \mathbf{n} ds = (dx_2, -dx_1).$$

67 Here t is the tangent vector along the boundary curve and ds is the element of arc-length. From (2.2) we get for $B \subset \mathbb{R}^2$ that

$$\begin{aligned} & \int_B \frac{\partial F_1(x_1, x_2)}{\partial x_1} dx_1 dx_2 + \int_B \frac{\partial F_2(x_1, x_2)}{\partial x_2} dx_1 dx_2 \\ &= \int_{\partial B} F_1(x_1, x_2) dx_2 - \int_{\partial B} F_2(x_1, x_2) dx_1, \end{aligned} \quad (2.3)$$

68 which is Green's theorem in plane.

If $n = 3$ and if ∂B is described as a level-set of a function of 3 variables i.e., $\partial B = \{x_1, x_2, x_3 \in \mathbb{R}^3 | G(x_1, x_2, x_3) = 0\}$, then a vector pointing in the direction of \mathbf{n} is $\text{grad}G$. We shall use the case where $G(x_1, x_2, x_3) = x_3 - g(x_1, x_2)$, $(x_1, x_2) \in D$, a domain in \mathbb{R}^2 for some differentiable function g on D and B corresponds to the inequality $x_3 < g(x_1, x_2)$, namely

$$B = \{(x_1, x_2, x_3) \in \mathbb{R}^3 | x_3 < g(x_1, x_2)\}.$$

Then

$$\mathbf{n} = \frac{(-g_{x_1}, -g_{x_2}, 1)}{(1 + g_{x_1}^2 + g_{x_2}^2)^{1/2}}, \quad dA = (1 + g_{x_1}^2 + g_{x_2}^2)^{1/2} dx_1 dx_2$$

and

$$\mathbf{n} dA = (-g_{x_1}, -g_{x_2}, 1) dx_1 dx_2.$$

From (2.2) we get

$$\begin{aligned} & \int_B \left(\frac{\partial F_1(x_1, x_2, x_3)}{\partial x_1} + \frac{\partial F_2(x_1, x_2, x_3)}{\partial x_2} + \frac{\partial F_3(x_1, x_2, x_3)}{\partial x_3} \right) dx_1 dx_2 dx_3 \\ &= - \int_D F_1(x_1, x_2, g(x_1, x_2)) g_{x_1}(x_1, x_2) dx_1 dx_2 \\ & \quad - \int_D F_2(x_1, x_2, g(x_1, x_2)) g_{x_2}(x_1, x_2) dx_1 dx_2 \\ & \quad + \int_D F_3(x_1, x_2, g(x_1, x_2)) dx_1 dx_2, \end{aligned} \quad (2.4)$$

69 which is the Gauss-Ostrogradsky theorem in space.

70 Following Apostol [1], we can also consider a surface described by the vector equation

$$r(u, v) = x_1(u, v) \vec{i} + x_2(u, v) \vec{j} + x_3(u, v) \vec{k}, \quad (2.5)$$

71 where $(u, v) \in [a, b] \times [c, d]$.

If x_1, x_2, x_3 are differentiable on $[a, b] \times [c, d]$ we consider the two vectors

$$\begin{aligned}\frac{\partial r}{\partial u} &= \frac{\partial x_1}{\partial u} \vec{i} + \frac{\partial x_2}{\partial u} \vec{j} + \frac{\partial x_3}{\partial u} \vec{k}, \\ \frac{\partial r}{\partial v} &= \frac{\partial x_1}{\partial v} \vec{i} + \frac{\partial x_2}{\partial v} \vec{j} + \frac{\partial x_3}{\partial v} \vec{k}.\end{aligned}$$

The cross product of these two vectors $\frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v}$ will be referred to as the fundamental vector product of the representation r . Its components can be expressed as Jacobian determinants. In fact, we have [1, p. 420]

$$\begin{aligned}\frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} &= \begin{vmatrix} \frac{\partial x_2}{\partial u} & \frac{\partial x_3}{\partial u} \\ \frac{\partial x_2}{\partial v} & \frac{\partial x_3}{\partial v} \end{vmatrix} \vec{i} + \begin{vmatrix} \frac{\partial x_3}{\partial u} & \frac{\partial x_1}{\partial u} \\ \frac{\partial x_3}{\partial v} & \frac{\partial x_1}{\partial v} \end{vmatrix} \vec{j} + \begin{vmatrix} \frac{\partial x_1}{\partial u} & \frac{\partial x_2}{\partial u} \\ \frac{\partial x_1}{\partial v} & \frac{\partial x_2}{\partial v} \end{vmatrix} \vec{k} \\ &= \frac{\partial(x_2, x_3)}{\partial(u, v)} \vec{i} + \frac{\partial(x_3, x_1)}{\partial(u, v)} \vec{j} + \frac{\partial(x_1, x_2)}{\partial(u, v)} \vec{k}.\end{aligned}\quad (2.6)$$

Let $\partial B = r(T)$ be a parametric surface described by a vector-valued function r defined on the box $T = [a, b] \times [c, d]$. The area of ∂B denoted $A_{\partial B}$ is defined by the double integral [1, pp. 424-425]

$$\begin{aligned}A_{\partial B} &= \int_a^b \int_c^d \left\| \frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} \right\| dudv \\ &= \int_a^b \int_c^d \sqrt{\left(\frac{\partial(x_2, x_3)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(x_3, x_1)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(x_1, x_2)}{\partial(u, v)} \right)^2} dudv.\end{aligned}\quad (2.7)$$

72 We define surface integrals in terms of a parametric representation for the surface. One
73 can prove that under certain general conditions the value of the integral is independent
74 of the representation.

Let $\partial B = r(T)$ be a parametric surface described by a vector-valued differentiable function r defined on the box $T = [a, b] \times [c, d]$ and let $f : \partial B \rightarrow \mathbb{C}$ defined and bounded on ∂B . The surface integral of f over ∂B is defined by [1, p. 430]

$$\begin{aligned}\iint_{\partial B} f dA &= \int_a^b \int_c^d f(x_1, x_2, x_3) \left\| \frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} \right\| dudv \\ &= \int_a^b \int_c^d f(x_1(u, v), x_2(u, v), x_3(u, v)) \\ &\quad \times \sqrt{\left(\frac{\partial(x_2, x_3)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(x_3, x_1)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(x_1, x_2)}{\partial(u, v)} \right)^2} dudv.\end{aligned}\quad (2.8)$$

If $\partial B = r(T)$ is a parametric surface, the fundamental vector product $N = \frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v}$ is normal to ∂B at each regular point of the surface. At each such point there are two unit

normals, a unit normal \mathbf{n}_1 , which has the same direction as N , and a unit normal \mathbf{n}_2 which has the opposite direction. Thus

$$\mathbf{n}_1 = \frac{N}{\|N\|} \quad \text{and} \quad \mathbf{n}_2 = -\mathbf{n}_1.$$

Let \mathbf{n} be one of the two normals \mathbf{n}_1 or \mathbf{n}_2 . Let also F be a vector field defined on ∂B and assume that the surface integral,

$$\iint_{\partial B} (F \cdot \mathbf{n}) dA,$$

75 called the flux surface integral, exists. Here $F \cdot \mathbf{n}$ is the dot or inner product.

We can write [1, p. 434]

$$\iint_{\partial B} (F \cdot \mathbf{n}) dA = \pm \int_a^b \int_c^d F(r(u, v)) \cdot \left(\frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} \right) dudv,$$

76 where the sign "+" is used if $\mathbf{n} = \mathbf{n}_1$ and the "-" sign is used if $\mathbf{n} = \mathbf{n}_2$.

If

$$\begin{aligned} F(x_1, x_2, x_3) &= F_1(x_1, x_2, x_3) \vec{i} + F_2(x_1, x_2, x_3) \vec{j} + F_3(x_1, x_2, x_3) \vec{k}, \\ r(u, v) &= x_1(u, v) \vec{i} + x_2(u, v) \vec{j} + x_3(u, v) \vec{k}, \quad \text{where } (u, v) \in [a, b] \times [c, d], \end{aligned}$$

then the flux surface integral for $\mathbf{n} = \mathbf{n}_1$ can be explicitly calculated as [1, p. 435]

$$\begin{aligned} \iint_{\partial B} (F \cdot \mathbf{n}) dA &= \int_a^b \int_c^d F_1(x_1(u, v), x_2(u, v), x_3(u, v)) \frac{\partial(x_2, x_3)}{\partial(u, v)} dudv \\ &\quad + \int_a^b \int_c^d F_2(x_1(u, v), x_2(u, v), x_3(u, v)) \frac{\partial(x_3, x_1)}{\partial(u, v)} dudv \\ &\quad + \int_a^b \int_c^d F_3(x_1(u, v), x_2(u, v), x_3(u, v)) \frac{\partial(x_1, x_2)}{\partial(u, v)} dudv. \end{aligned} \quad (2.9)$$

The sum of the double integrals on the right is often written more briefly as [1, p. 435]

$$\begin{aligned} \iint_{\partial B} F_1(x_1, x_2, x_3) dx_2 \wedge dx_3 + \iint_{\partial B} F_2(x_1, x_2, x_3) dx_3 \wedge dx_1 \\ + \iint_{\partial B} F_3(x_1, x_2, x_3) dx_1 \wedge dx_2. \end{aligned}$$

77 Let $B \subset \mathbb{R}^3$ be a solid in 3-space bounded by an orientable closed surface ∂B , and let

78 \mathbf{n} be the unit outer normal to ∂B . If F is a continuously differentiable vector field defined

79 on B , we have the Gauss-Ostrogradsky identity

$$\iiint_B (\operatorname{div} F) dV = \iint_{\partial B} (F \cdot \mathbf{n}) dA. \quad (2.10)$$

If we express

$$F(x_1, x_2, x_3) = F_1(x_1, x_2, x_3) \vec{i} + F_2(x_1, x_2, x_3) \vec{j} + F_3(x_1, x_2, x_3) \vec{k},$$

then (2.4) can be written as

$$\begin{aligned} & \iiint_B \left(\frac{\partial F_1(x_1, x_2, x_3)}{\partial x_1} + \frac{\partial F_2(x_1, x_2, x_3)}{\partial x_2} + \frac{\partial F_3(x_1, x_2, x_3)}{\partial x_3} \right) dx_1 dx_2 dx_3 \\ &= \iint_{\partial B} F_1(x_1, x_2, x_3) dx_2 \wedge dx_3 + \iint_{\partial B} F_2(x_1, x_2, x_3) dx_3 \wedge dx_1 \\ & \quad + \iint_{\partial B} F_3(x_1, x_2, x_3) dx_1 \wedge dx_2. \end{aligned} \quad (2.11)$$

80 3 Main results

81 We start with the following identity that is of interest in itself:

Lemma 3.1. Assume that $f : D \rightarrow \mathbb{C}$ has partial derivatives on the domain $D \subset \mathbb{R}^n$, $n \geq 2$. Define for $j \neq i$

$$\Lambda_{\partial f, D}(x_i, x_j) := (x_i - x_j) \left(\frac{\partial f(x_1, \dots, x_n)}{\partial x_i} - \frac{\partial f(x_1, \dots, x_n)}{\partial x_j} \right),$$

where $(x_1, \dots, x_n) \in D$. Then we have

$$\begin{aligned} & \frac{1}{n-1} \sum_{k=1}^n \frac{\partial}{\partial x_k} \left(\left(x_k - \frac{1}{n} \sum_{j=1}^n x_j \right) f(x_1, \dots, x_n) \right) \\ &= f(x_1, \dots, x_n) + \frac{1}{n(n-1)} \sum_{1 \leq i < j \leq n} \Lambda_{\partial f, D}(x_i, x_j). \end{aligned} \quad (3.1)$$

Proof. For $j \neq i$ we have

$$\begin{aligned} \frac{\partial}{\partial x_i} ((x_i - x_j) f(x_1, \dots, x_n)) &= f(x_1, \dots, x_n) + (x_i - x_j) \frac{\partial f(x_1, \dots, x_n)}{\partial x_i}, \\ \frac{\partial}{\partial x_j} ((x_i - x_j) f(x_1, \dots, x_n)) &= -f(x_1, \dots, x_n) + (x_i - x_j) \frac{\partial f(x_1, \dots, x_n)}{\partial x_j}, \end{aligned}$$

which gives

$$\begin{aligned} & \frac{\partial}{\partial x_i} ((x_i - x_j) f(x_1, \dots, x_n)) - \frac{\partial}{\partial x_j} ((x_i - x_j) f(x_1, \dots, x_n)) \\ &= 2f(x_1, \dots, x_n) + (x_i - x_j) \left(\frac{\partial f(x_1, \dots, x_n)}{\partial x_i} - \frac{\partial f(x_1, \dots, x_n)}{\partial x_j} \right) \end{aligned}$$

82 for $j \neq i$.

If we take the sum over $i, j \in \{1, \dots, n\}$ with $j \neq i$ we get

$$\begin{aligned} & \sum_{i,j=1,j \neq i}^n \left[\frac{\partial}{\partial x_i} ((x_i - x_j) f(x_1, \dots, x_n)) - \frac{\partial}{\partial x_j} ((x_i - x_j) f(x_1, \dots, x_n)) \right] \\ &= 2 \sum_{i,j=1,j \neq i}^n f(x_1, \dots, x_n) + \sum_{i,j=1,j \neq i}^n (x_i - x_j) \left(\frac{\partial f(x_1, \dots, x_n)}{\partial x_i} - \frac{\partial f(x_1, \dots, x_n)}{\partial x_j} \right). \quad (3.2) \end{aligned}$$

We have

$$\begin{aligned} & \sum_{i,j=1,j \neq i}^n f(x_1, \dots, x_n) = n(n-1) f(x_1, \dots, x_n), \\ & \sum_{i,j=1,j \neq i}^n (x_i - x_j) \left(\frac{\partial f(x_1, \dots, x_n)}{\partial x_i} - \frac{\partial f(x_1, \dots, x_n)}{\partial x_j} \right) \\ &= 2 \sum_{1 \leq i < j \leq n} (x_i - x_j) \left(\frac{\partial f(x_1, \dots, x_n)}{\partial x_i} - \frac{\partial f(x_1, \dots, x_n)}{\partial x_j} \right). \end{aligned}$$

Also

$$\begin{aligned} & \sum_{i,j=1,j \neq i}^n \left[\frac{\partial}{\partial x_i} ((x_i - x_j) f(x_1, \dots, x_n)) - \frac{\partial}{\partial x_j} ((x_i - x_j) f(x_1, \dots, x_n)) \right] \\ &= \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\sum_{j=1,j \neq i}^n (x_i - x_j) f(x_1, \dots, x_n) \right) \\ & \quad - \sum_{j=1}^n \frac{\partial}{\partial x_j} \left(\sum_{i=1,j \neq i}^n (x_i - x_j) f(x_1, \dots, x_n) \right) \\ &= \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\left((n-1)x_i - \sum_{j=1,j \neq i}^n x_j \right) f(x_1, \dots, x_n) \right) \\ & \quad - \sum_{j=1}^n \frac{\partial}{\partial x_j} \left(\left(\sum_{i=1,j \neq i}^n x_i - (n-1)x_j \right) f(x_1, \dots, x_n) \right) \\ &= \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\left((n-1)x_i - \sum_{j=1,j \neq i}^n x_j \right) f(x_1, \dots, x_n) \right) \\ & \quad + \sum_{j=1}^n \frac{\partial}{\partial x_j} \left(\left((n-1)x_j - \sum_{i=1,j \neq i}^n x_i \right) f(x_1, \dots, x_n) \right) \\ &= 2 \sum_{k=1}^n \frac{\partial}{\partial x_k} \left(\left((n-1)x_k - \sum_{j=1,j \neq k}^n x_j \right) f(x_1, \dots, x_n) \right) \end{aligned}$$

$$= 2 \sum_{k=1}^n \frac{\partial}{\partial x_k} \left(\left(nx_k - \sum_{j=1}^n x_j \right) f(x_1, \dots, x_n) \right).$$

By (3.2) we get

$$\begin{aligned} & 2 \sum_{k=1}^n \frac{\partial}{\partial x_k} \left(\left(nx_k - \sum_{j=1}^n x_j \right) f(x_1, \dots, x_n) \right) \\ &= 2n(n-1) f(x_1, \dots, x_n) \\ & \quad + 2 \sum_{1 \leq i < j \leq n} (x_i - x_j) \left(\frac{\partial f(x_1, \dots, x_n)}{\partial x_i} - \frac{\partial f(x_1, \dots, x_n)}{\partial x_j} \right), \end{aligned}$$

83 which is equivalent to the desired result. □

Remark 3.1. For $n = 2$ we get

$$\begin{aligned} & \frac{1}{2} \left[\frac{\partial}{\partial x_1} [(x_1 - x_2) f(x_1, x_2)] + \frac{\partial}{\partial x_1} [(x_2 - x_1) f(x_1, x_2)] \right] \\ &= f(x_1, x_2) + \frac{1}{2} \Lambda_{\partial f, D}(x_1, x_2) \end{aligned} \quad (3.3)$$

for $(x_1, x_2) \in D$. For $n = 3$ we get

$$\begin{aligned} & \frac{1}{3} \left[\frac{\partial}{\partial x_1} \left(\left(x_1 - \frac{x_2 + x_3}{2} \right) f(x_1, x_2, x_3) \right) + \frac{\partial}{\partial x_2} \left(\left(x_2 - \frac{x_1 + x_3}{2} \right) f(x_1, x_2, x_3) \right) \right. \\ & \quad \left. + \frac{\partial}{\partial x_3} \left(\left(x_3 - \frac{x_1 + x_2}{2} \right) f(x_1, x_2, x_3) \right) \right] \\ &= f(x_1, x_2, x_3) + \frac{1}{6} [\Lambda_{\partial f, D}(x_1, x_2) + \Lambda_{\partial f, D}(x_2, x_3) + \Lambda_{\partial f, D}(x_1, x_3)] \end{aligned} \quad (3.4)$$

84 for $(x_1, x_2, x_3) \in D$.

85 We have the following identity of interest:

Theorem 3.1. Let B be a bounded closed subset of \mathbb{R}^n ($n \geq 2$) with smooth (or piecewise smooth) boundary ∂B and $\mathbf{n} = (\mathbf{n}_1, \dots, \mathbf{n}_n)$ be the unit outward-pointing normal of ∂B . If f is a continuously differentiable function on an open neighborhood of B , then we have the representation

$$\begin{aligned} & \frac{1}{n-1} \sum_{k=1}^n \int_{\partial B} \left(x_k - \frac{1}{n} \sum_{j=1}^n x_j \right) f(x) \mathbf{n}_k(x) dA - \int_B f(x) dx \\ &= \frac{1}{n(n-1)} \sum_{1 \leq i < j \leq n} \int_B \Lambda_{\partial f, B}(x_i, x_j) dx. \end{aligned} \quad (3.5)$$

Proof. We use the identity (3.1) on B for $x = (x_1, \dots, x_n)$ and take the volume integral to get

$$\begin{aligned} & \frac{1}{n-1} \int_B \sum_{k=1}^n \frac{\partial}{\partial x_k} \left(\left(x_k - \frac{1}{n} \sum_{j=1}^n x_j \right) f(x) \right) dx \\ &= \int_B f(x) dx + \frac{1}{n(n-1)} \sum_{1 \leq i < j \leq n} \int_B \Lambda_{\partial f, B}(x_i, x_j) dx. \end{aligned} \quad (3.6)$$

Define

$$F_k(x) = \left(x_k - \frac{1}{n} \sum_{j=1}^n x_j \right) f(x), \quad k \in \{1, \dots, n\}, \quad x \in B,$$

and use the Divergence theorem (2.2) to get

$$\begin{aligned} & \int_B \sum_{k=1}^n \frac{\partial}{\partial x_k} \left(\left(x_k - \frac{1}{n} \sum_{j=1}^n x_j \right) f(x) \right) dx \\ &= \sum_{k=1}^n \int_{\partial B} \left(x_k - \frac{1}{n} \sum_{j=1}^n x_j \right) f(x) \mathbf{n}_k(x) dA. \end{aligned} \quad (3.7)$$

On utilising (3.6) and (3.7), we obtain

$$\begin{aligned} & \int_B f(x) dx + \frac{1}{n(n-1)} \sum_{1 \leq i < j \leq n} \int_B \Lambda_{\partial f, B}(x_i, x_j) dx \\ &= \frac{1}{n-1} \sum_{k=1}^n \int_{\partial B} \left(x_k - \frac{1}{n} \sum_{j=1}^n x_j \right) f(x) \mathbf{n}_k(x) dA, \end{aligned}$$

⁸⁶ that is equivalent to (3.5). □

Remark 3.2. For $n = 2$ we obtain the identity

$$\begin{aligned} & \frac{1}{2} \int_{\partial B} [(x_1 - x_2) f(x_1, x_2) dx_1 + (x_1 - x_2) f(x_1, x_2) dx_2] \\ & \quad - \int_B f(x_1, x_2) dx_1 dx_2 \\ &= \frac{1}{2} \int_B \Lambda_{\partial f, B}(x_1, x_2) dx_1 dx_2, \end{aligned} \quad (3.8)$$

⁸⁷ where B is a bounded closed subset of \mathbb{R}^2 with smooth (or piecewise smooth) boundary
⁸⁸ ∂B and f is a continuously differentiable function on an open neighborhood of B .

For $n = 3$ we obtain the identity

$$\begin{aligned} & \frac{1}{3} \left[\int_{\partial B} \left(x_1 - \frac{x_2 + x_3}{2} \right) f(x_1, x_2, x_3) dx_2 \wedge dx_3 \right. \\ & \quad + \int_{\partial B} \left(x_2 - \frac{x_1 + x_3}{2} \right) f(x_1, x_2, x_3) dx_3 \wedge dx_1 \\ & \quad \left. + \int_{\partial B} \left(x_3 - \frac{x_1 + x_2}{2} \right) f(x_1, x_2, x_3) dx_1 \wedge dx_2 \right] - \int_B f(x_1, x_2, x_3) dx_1 dx_2 dx_3 \\ & = \frac{1}{6} \int_B [\Lambda_{\partial f, B}(x_1, x_2) + \Lambda_{\partial f, B}(x_2, x_3) + \Lambda_{\partial f, B}(x_1, x_3)] dx_1 dx_2 dx_3, \end{aligned}$$

89 where B is a bounded closed subset of \mathbb{R}^3 with smooth (or piecewise smooth) boundary
90 ∂B and f is a continuously differentiable function on an open neighborhood of B .

91 **Corollary 3.1.** Let B be a bounded closed and symmetric convex subset of \mathbb{R}^n ($n \geq 2$) with
92 smooth (or piecewise smooth) boundary ∂B and $\mathbf{n} = (\mathbf{n}_1, \dots, \mathbf{n}_n)$ be the unit outward-pointing
93 normal of ∂B . If f is a continuously differentiable function on an open neighborhood of B and
94 Schur convex on B , then we have the integral inequality

$$\frac{1}{n-1} \sum_{k=1}^n \int_{\partial B} \left(x_k - \frac{1}{n} \sum_{j=1}^n x_j \right) f(x) \mathbf{n}_k(x) dA \geq \int_B f(x) dx. \quad (3.9)$$

95 *Proof.* Since f is Schur convex on B , then by (1.3) we get $\Lambda_{\partial f, D}(x_i, x_j) \geq 0$ for all $1 \leq i <$
96 $j \leq n$, and by using (3.5) we get the desired inequality (3.9). \square

97 **Corollary 3.2.** With the assumptions of Corollary 3.1 and if there exists $L_{ij} > 0$ for $1 \leq i < j \leq$
98 n such that

$$\Lambda_{\partial f, D}(x_i, x_j) \leq L_{ij} (x_i - x_j)^2 \quad \text{for all } x = (x_1, \dots, x_n) \in B, \quad (3.10)$$

then we also have the reverse inequality

$$\begin{aligned} 0 & \leq \frac{1}{n-1} \sum_{k=1}^n \int_{\partial B} \left(x_k - \frac{1}{n} \sum_{j=1}^n x_j \right) f(x) \mathbf{n}_k(x) dA - \int_B f(x) dx \\ & \leq \frac{1}{n(n-1)} \sum_{1 \leq i < j \leq n} L_{ij} \int_B (x_i - x_j)^2 dx. \end{aligned} \quad (3.11)$$

99 The proof follows by the equality (3.5).

Remark 3.3. For $n = 2$ in (3.9) we get

$$\begin{aligned} 0 & \leq \frac{1}{2} \int_{\partial B} [(x_1 - x_2) f(x_1, x_2) dx_1 + (x_1 - x_2) f(x_1, x_2) dx_2] \\ & \quad - \int_B f(x_1, x_2) dx_1 dx_2 \\ & \leq \frac{1}{2} L \int_B (x_1 - x_2)^2 dx_1 dx_2, \end{aligned} \quad (3.12)$$

provided that f is Schur convex on the convex and symmetric domain $B \subset \mathbb{R}^2$ and there exists $L > 0$ such that

$$\begin{aligned} \Lambda_{\partial f, D}(x_1, x_2) &= (x_1 - x_2) \left(\frac{\partial f(x_1, x_2)}{\partial x_1} - \frac{\partial f(x_1, x_2)}{\partial x_2} \right) \\ &\leq L(x_1 - x_2)^2 \quad \text{for all } x = (x_1, x_2) \in B. \end{aligned} \quad (3.13)$$

For $n = 3$ we get

$$0 \leq \frac{1}{3} \left[\int_{\partial B} \left(x_1 - \frac{x_2 + x_3}{2} \right) f(x_1, x_2, x_3) dx_2 \wedge dx_3 \right. \quad (3.14)$$

$$+ \int_{\partial B} \left(x_2 - \frac{x_1 + x_3}{2} \right) f(x_1, x_2, x_3) dx_3 \wedge dx_1$$

$$\left. + \int_{\partial B} \left(x_3 - \frac{x_1 + x_2}{2} \right) f(x_1, x_2, x_3) dx_1 \wedge dx_2 \right]$$

$$- \int_B f(x_1, x_2, x_3) dx_1 dx_2 dx_3$$

$$\leq \frac{1}{6} \left[L_{12} \int_B (x_1 - x_2)^2 dx_1 dx_2 dx_3 \right.$$

$$\left. + L_{23} \int_B (x_2 - x_3)^2 dx_1 dx_2 dx_3 + L_{13} \int_B (x_1 - x_3)^2 dx_1 dx_2 dx_3 \right] \quad (3.15)$$

provided that f is Schur convex on the convex and symmetric domain $B \subset \mathbb{R}^3$ and

$$\begin{aligned} \Lambda_{\partial f, D}(x_i, x_j) &= (x_i - x_j) \left(\frac{\partial f(x_1, x_2, x_3)}{\partial x_i} - \frac{\partial f(x_1, x_2, x_3)}{\partial x_j} \right) \\ &\leq L_{ij} (x_i - x_j)^2 \quad \text{for all } x = (x_1, x_2, x_3) \in B, \end{aligned} \quad (3.16)$$

100 where $L_{ij} > 0$ for $1 \leq i < j \leq 3$.

101 4 An example for three dimensional balls

Consider the 3-dimensional ball centered in $O = (0, 0, 0)$ and having the radius $R > 0$,

$$B(O, R) := \{ (x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_1^2 + x_2^2 + x_3^2 \leq R^2 \}$$

and the sphere

$$S(O, R) := \{ (x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_1^2 + x_2^2 + x_3^2 = R^2 \}.$$

Consider the parametrization of $B(O, R)$ and $S(O, R)$ given by:

$$B(O, R) : \begin{cases} x_1 = r \cos \psi \cos \varphi, \\ x_2 = r \cos \psi \sin \varphi, \\ x_3 = r \sin \psi, \end{cases} \quad (r, \psi, \varphi) \in [0, R] \times \left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \times [0, 2\pi],$$

and

$$S(O, R) : \begin{cases} x_1 = R \cos \psi \cos \varphi, \\ x_2 = R \cos \psi \sin \varphi, \\ x_3 = R \sin \psi, \end{cases} \quad (\psi, \varphi) \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \times [0, 2\pi].$$

We have

$$\begin{aligned} \begin{vmatrix} \frac{\partial x_2}{\partial \psi} & \frac{\partial x_3}{\partial \psi} \\ \frac{\partial x_2}{\partial \varphi} & \frac{\partial x_3}{\partial \varphi} \end{vmatrix} &= -R^2 \cos^2 \psi \cos \varphi, & \begin{vmatrix} \frac{\partial x_1}{\partial \psi} & \frac{\partial x_3}{\partial \psi} \\ \frac{\partial x_1}{\partial \varphi} & \frac{\partial x_3}{\partial \varphi} \end{vmatrix} &= R^2 \cos^2 \psi \sin \varphi, \\ \begin{vmatrix} \frac{\partial x_1}{\partial \psi} & \frac{\partial x_2}{\partial \psi} \\ \frac{\partial x_1}{\partial \varphi} & \frac{\partial x_2}{\partial \varphi} \end{vmatrix} &= -R^2 \sin \psi \cos \psi. \end{aligned}$$

In Cartesian coordinates, we have the inequality (3.14) written as

$$\begin{aligned} 0 &\leq \frac{1}{3} \left[\int_{S(O,R)} \left(x_1 - \frac{x_2 + x_3}{2} \right) f(x_1, x_2, x_3) dx_2 \wedge dx_3 \right. \\ &\quad + \int_{S(O,R)} \left(x_2 - \frac{x_1 + x_3}{2} \right) f(x_1, x_2, x_3) dx_3 \wedge dx_1 \\ &\quad + \int_{S(O,R)} \left(x_3 - \frac{x_1 + x_2}{2} \right) f(x_1, x_2, x_3) dx_1 \wedge dx_2 \left. \right] \\ &\quad - \int_{B(O,R)} f(x_1, x_2, x_3) dx_1 dx_2 dx_3 \\ &\leq \frac{1}{6} \left[L_{12} \int_{B(O,R)} (x_1 - x_2)^2 dx_1 dx_2 dx_3 \right. \\ &\quad \left. + L_{23} \int_{B(O,R)} (x_2 - x_3)^2 dx_1 dx_2 dx_3 + L_{13} \int_{B(O,R)} (x_1 - x_3)^2 dx_1 dx_2 dx_3 \right] \quad (4.1) \end{aligned}$$

102 provided that f is a continuously differentiable function on an open neighborhood of
103 $B(O, R)$, Schur convex on $B(O, R)$ and the condition (3.16) is fulfilled.

Now, observe that

$$\begin{aligned} &\int_{B(O,R)} (x_1 - x_2)^2 dx_1 dx_2 dx_3 \\ &= \int_0^R \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2\pi} (r \cos \psi \cos \varphi - r \cos \psi \sin \varphi)^2 r^2 \cos \psi dr d\psi d\varphi \\ &= \int_0^R r^4 dr \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^3 \psi d\psi \int_0^{2\pi} (\cos \varphi - \sin \varphi)^2 d\varphi = \frac{R^5}{5} \left(\frac{4}{3} \right) 2\pi \\ &= \frac{8}{15} \pi R^5 \end{aligned}$$

and, similarly

$$\int_{B(O,R)} (x_2 - x_3)^2 dx_1 dx_2 dx_3 = \int_{B(O,R)} (x_1 - x_3)^2 dx_1 dx_2 dx_3 = \frac{8}{15} \pi R^5.$$

In polar coordinates, (4.1) becomes

$$\begin{aligned} 0 \leq & \frac{1}{3} R^3 \left[- \int_{S(O,R)} \left(\cos \psi \cos \varphi - \frac{\cos \psi \sin \varphi + \sin \psi}{2} \right) \right. \\ & \times f(R \cos \psi \cos \varphi, R \cos \psi \sin \varphi, R \sin \psi) \cos^2 \psi \cos \varphi d\psi d\varphi \\ & + \int_{S(O,R)} \left(\cos \psi \sin \varphi - \frac{\cos \psi \cos \varphi + \sin \psi}{2} \right) \\ & \times f(R \cos \psi \cos \varphi, R \cos \psi \sin \varphi, R \sin \psi) \cos^2 \psi \sin \varphi d\psi d\varphi \\ & - \int_{S(O,R)} \left(\sin \psi - \frac{\cos \psi \cos \varphi + \cos \psi \sin \varphi}{2} \right) \\ & \times f(R \cos \psi \cos \varphi, R \cos \psi \sin \varphi, R \sin \psi) \sin \psi \cos \varphi d\psi d\varphi \\ & \left. - \int_0^R \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2\pi} f(r \cos \psi \cos \varphi, r \cos \psi \sin \varphi, r \sin \psi) r^2 \cos \psi dr d\psi d\varphi \right] \\ \leq & \frac{4}{45} \pi R^5 (L_{12} + L_{23} + L_{13}), \end{aligned} \quad (4.2)$$

104 provided that f is a continuously differentiable function on an open neighborhood of
105 $B(O, R)$, Schur convex on $B(O, R)$ and satisfying the condition (3.16).

106 References

- 107 [1] T. M. Apostol, *Calculus Volume II, Multi Variable Calculus and Linear Algebra, with Appli-*
108 *cations to Differential Equations and Probability, Second Edition, John Wiley & Sons, New*
109 *York London Sydney Toronto, 1969.*
- 110 [2] V. Čuljak, A remark on Schur-convexity of the mean of a convex function. *J. Math. Inequal.*,
111 9(4) (2015), 1133–1142.
- 112 [3] S. S. Dragomir, Inequalities for double integrals of Schur convex functions on symmetric and
113 convex domains, *Mat. Vesnik*, 73(1) (2021), 63–74. Preprint RGMIA Research Rep. Coll., 22
114 (2019), <http://rgmia.org/papers/v22/v22a69.pdf>.
- 115 [4] S. S. Dragomir and K. Nikodem, Functions generating (m, M, Ψ) -Schur-convex sums, *Ae-*
116 *quationes Math.*, 93(1) (2019), 79–90.
- 117 [5] A. W. Marshall, I. Olkin and B. C. Arnold, *Inequalities: Theory of Majorization and Its Ap-*
118 *plications, Second Edition, Springer, 2011.*
- 119 [6] K. Nikodem, T. Rajba and S. Wasowicz, Functions generating strongly Schur-convex
120 sums, *Inequalities and applications 2010*, 175–182, *Int. Ser. Numer. Math.*, 161,
121 Birkhäuser/Springer, Basel, 2012.
- 122 [7] J. Qi and W. Wang, Schur convex functions and the Bonnesen style isoperimetric inequalities
123 for planar convex polygons. *J. Math. Inequal.*, 12(1) (2018), 23–29.

- 124 [8] H.-N. Shi and J. Zhang, Compositions involving Schur harmonically convex functions. J.
125 Comput. Anal. Appl., 22(5) (2017), 907–922.
- 126 [9] M. Singer, The divergence theorem, [https://www.maths.ed.ac.uk/~jmf/Teaching/](https://www.maths.ed.ac.uk/~jmf/Teaching/Lectures/divthm.pdf)
127 Lectures/divthm.pdf.