

On Symmetry Breaking of Allen-Cahn

Dong Li^{1,2}, Chaoyu Quan^{1,*}, Tao Tang^{3,4} and Wen Yang⁵

¹ *SUSTech International Center for Mathematics, Southern University of Science and Technology, Shenzhen, P.R. China.*

² *Department of Mathematics, Southern University of Science and Technology, Shenzhen, P.R. China.*

³ *Guangdong Provincial Key Laboratory of Computational Science and Material Design, Southern University of Science and Technology, Shenzhen, P.R. China.*

⁴ *Division of Science and Technology, BNU-HKBU United International College, Zhuhai, P.R. China.*

⁵ *Wuhan Institute of Physics and Mathematics, Innovation Academy for Precision Measurement Science and Technology, Chinese Academy of Sciences, P.O. Box 71010, Wuhan 430071, P.R. China.*

Received 17 May 2021; Accepted 20 July 2021

Abstract. This paper is concerned with numerical solutions for the Allen-Cahn equation with standard double well potential and periodic boundary conditions. Surprisingly it is found that using standard numerical discretizations with high precision computational solutions may converge to completely incorrect steady states. This happens for very smooth initial data and state-of-the-art algorithms. We analyze this phenomenon and showcase the resolution of this problem by a new symmetry-preserving filter technique. We develop a new theoretical framework and rigorously prove the convergence to steady states for the filtered solutions.

AMS subject classifications: 35K55, 65M12

Key words: Allen-Cahn equation, symmetry breaking, steady state.

1 Introduction

A curious experiment. Consider the following 1D Allen-Cahn on the periodic torus $\mathbb{T} = [-\pi, \pi]$:

$$\begin{cases} \partial_t u = \kappa^2 \partial_{xx} u - f(u), \\ u|_{t=0} = u_0, \end{cases} \quad (1.1)$$

*Corresponding author. *Email addresses:* lid@sustech.edu.cn (D. Li), quancy@sustech.edu.cn (C. Quan), ttang@uic.edu.cn (T. Tang), wyang@wipm.ac.cn (W. Yang)




Figure 1: Dynamics of the Allen-Cahn equation using the first-order implicit-explicit scheme with different discretization parameters. (a): $\kappa=1$, $N_p=128$ and $\tau=0.01$, and (b): $\kappa=1$, $N_p=4096$ and $\tau=10^{-5}$.

where $f(u) = u^3 - u$ corresponds to the usual double-well potential, $\kappa^2 > 0$ is the diffusion coefficient. The scalar function $u: \mathbb{T} \rightarrow \mathbb{R}$ represents the concentration of a phase in an alloy and typically has values in the physical range $[-1, 1]$. There is by now an extensive literature on the theoretical analysis and numerical simulation of the Allen-Cahn equation and related phase field models (cf. [1–5, 7, 8, 14]). For the system (1.1) we take the initial data u_0 as an odd function of x (here we tacitly “lift” the periodic function u to be defined on the whole real axis so that oddness can be defined in the usual way). For example, one can take $u_0(x) = \sin(x)$. Denote

$$u_\infty(x) = \lim_{t \rightarrow \infty} u(x, t). \quad (1.2)$$

Since u_0 is odd and smooth, it is clear that the odd symmetry should be preserved for all time. In particular, the final state u_∞ must be a periodic odd function of x . However to our surprise, standard numerical experiments show that this parity property may be lost in not very long time simulations. For example, by using the finite difference method with $N_p = 128$ nodes for space discretization and the first-order implicit-explicit method with time step $\tau=0.01$ for time discretization, we find that u tends to ± 1 when t is suitably large (see the left-hand side of Fig. 1). Obviously, ± 1 are not the correct steady states as the oddness is not preserved.

To check the fidelity of the numerical scheme and rule out the issues connected with inaccurate numerical discretization, we first test larger $N = 4096$ and smaller $\tau = 10^{-5}$. It turns out that for this case, the oddness is preserved up to around $t \approx 32$, after which the solution u lost its oddness apparently and tends to 1 quickly. See the right-hand side of Fig. 1.




Figure 2: Dynamics of the Allen-Cahn equation using different numerical schemes ($\kappa=1$, $N_p=128$, $\tau=0.01$): implicit Euler method [16], the first-order IMEX method, the second-order operator splitting methods [6], BDF2 extrapolation method [15], the first- and second-order scalar auxiliary variable methods [12].

How about other numerical schemes? We test different first and second order (in time) numerical schemes to find that all these methods suffer the same non-physical phenomenon, see Fig. 2. This appears to be quite a serious issue, since it shows that all these standard numerical implementations may not be credible in not very long time simulations.

Outline of the paper. The purpose of this work is to rectify this issue for some special initial conditions by introducing a symmetry-preserving filter. The importance of this filter is that it eliminates the aforementioned unwanted projections into the unstable directions at each iteration. In the simplest situation of odd initial data such as $\sin x$, we dynamically force all the Fourier coefficient corresponding to cosine Fourier modes of the numerical solution to remain strictly zero after each iteration. For initial data such as $\sin(Lx)$ with $L \geq 2$ an integer, one can check that the spectral gap L is preserved in time, and our proposed filter forces the Fourier coefficients to retain the spectral gap as well as the symmetry conditions. Furthermore, we rigorously prove that the filtered solution will converge to the true steady state.

The rest of this paper is organized as follows. In Section 2 we classify the steady states and analyze their profiles. In Sections 3 and 4 we analyze the symmetry breaking phenomenon and show how the filtering approach can resolve this issue. In Section 5, by

providing more numerical examples we discuss in more details about various scenarios where the parity can be preserved or lost when the filter is in effect or turned off. It will be demonstrated that although the algorithm and analysis are provided for 1D Allen-Cahn the effectiveness of the filtering technique is observed for the 2D Allen-Cahn equation and 2D Cahn-Hilliard equation. In the last section we give some concluding remarks.

2 Properties of steady states

We consider the steady state solution of (1.1) which satisfies

$$\kappa^2 u'' + u - u^3 = 0, \quad x \in \mathbb{T}. \quad (2.1)$$

We first discuss a special class of steady states, called **odd zero-up ground states**.

Definition 2.1 (Odd zero-up ground state). *For any $0 < \kappa < 1$, U_κ is an odd zero-up ground state of (2.1) if U_κ is odd, $U'_\kappa(x) > 0$ on $[0, \frac{\pi}{2})$, and $U'_\kappa(x) < 0$ on $(\frac{\pi}{2}, \pi]$.*

To fix the notation, let us recall the energy functional associated with (1.1):

$$E(u) = \int_{\mathbb{T}} \left(\frac{\kappa^2}{2} (\partial_x u)^2 + \frac{1}{4} (1 - u^2)^2 \right) dx. \quad (2.2)$$

Remark 2.1. By a result from [10], for each $0 < \kappa < 1$ and excluding the trivial states $u \equiv \pm 1$, U_κ has the lowest energy among the steady state solutions, namely,

$$E_\kappa^{(0)} = \inf_{u \in \mathcal{F}} E(u) = E(U_\kappa), \quad (2.3)$$

where the class \mathcal{F} of admissible functions is given by

$$\mathcal{F} = \{u \mid u \in H^1(\mathbb{T}) \text{ solves (2.1) and } |u| < 1 \text{ for } x \in \mathbb{T}\}. \quad (2.4)$$

Furthermore, if $v \in \mathcal{F}$ attains E_κ , then for some $x_0 \in \mathbb{T}$, we have $v(x) = \pm U_\kappa(x - x_0)$. Note that the constraint $|u| < 1$ in (2.4) is to exclude the trivial minimizers $u = \pm 1$. Another way is to impose symmetry. For example U_κ or $-U_\kappa$ are the only minimizers amongst all odd 2π -periodic functions.

It is possible to classify all steady states to (2.1) by using U_κ . For any $0 < \kappa < 1$, define $m_\kappa \geq 1$ as the unique integer such that

$$\frac{1}{m_\kappa + 1} \leq \kappa < \frac{1}{m_\kappa}. \quad (2.5)$$

Theorem 2.1 ([10]). *Let $0 < \kappa < 1$. If u is a bounded solution to (2.1), then only one of the following holds.*

(1) $u \equiv \pm 1$ or $u \equiv 0$.

(2) $u = \pm U_{j\kappa}(jx+c)$, for some integer $j \in \{1, 2, \dots, m_\kappa\}$ and some constant c . Here note that $0 < j\kappa < 1$ by (2.5) and $U_{j\kappa}(j\cdot)$ can be viewed as a rescaled function of $U_{j\kappa}(\cdot)$, where $u(\cdot) = U_{j\kappa}(\cdot)$ corresponds to the odd zero-up ground state of the equation

$$j^2 \kappa^2 u'' + u - u^3 = 0, \quad x \in \mathbb{T}. \tag{2.6}$$

Remark 2.2. For $\kappa \geq 1$, it can be shown (see [10]) that if u is a bounded steady state solution to (2.1), then $u \equiv 0$ or ± 1 .

Roughly speaking, the significance of the odd zero-up ground state U_κ introduced in Definition 2.1 is that, the set $\{U_\kappa\}_{0 < \kappa < 1}$ gives a clean description of all possible bounded steady states of (2.1). Indeed by Theorem 2.1, any other bounded steady state must either be 0, ± 1 or a rescaled, translated copy from the catalogue $\{U_\kappa\}_{0 < \kappa < 1}$. Therefore it is of fundamental importance to study the odd zero-up ground states $\{U_\kappa\}_{0 < \kappa < 1}$.

We now characterize the detailed profile of the odd zero-up ground state of (2.1). Multiplying (2.1) (now u is replaced by U_κ) by U'_κ , integrating the resulting equation from $\frac{\pi}{2}$ to x , and using $U'_\kappa(\frac{\pi}{2}) = 0$ (a fact from the definition 2.1), we have

$$(U'_\kappa)^2 = \frac{1}{2\kappa^2} \left((U_\kappa^2 - 1)^2 - (N_\kappa^2 - 1)^2 \right), \tag{2.7}$$

where $N_\kappa := U_\kappa(\frac{\pi}{2})$ is the maximum of U_κ .

By (2.2) and (2.7), the ground state energy $E_\kappa^{(0)} = E(U_\kappa)$ can be rewritten as

$$E_\kappa^{(0)} = \int_{\mathbb{T}} \left(\frac{1}{2} (U_\kappa^2(x) - 1)^2 - \frac{1}{4} (N_\kappa^2 - 1)^2 \right) dx. \tag{2.8}$$

Since U_κ is monotonically increasing on $(0, \frac{\pi}{2})$, we have

$$U'_\kappa(x) = \frac{1}{\sqrt{2}\kappa} \sqrt{(U_\kappa^2 - 1)^2 - (N_\kappa^2 - 1)^2}, \quad x \in (0, \pi/2), \tag{2.9}$$

with $U_\kappa(0) = 0$. This yields

$$g(N_\kappa) := \int_0^{N_\kappa} \frac{1}{\sqrt{(u^2 - 1)^2 - (1 - N_\kappa^2)^2}} du = \int_0^{\frac{\pi}{2}} \frac{1}{\sqrt{2 - N_\kappa^2(1 + \sin^2 \theta)}} d\theta = \frac{\pi}{2\sqrt{2}\kappa}. \tag{2.10}$$

Note that $g(N_\kappa)$ is monotonically increasing on $[0, 1)$,

$$g(0) = \frac{\pi}{2\sqrt{2}}, \quad g(N_\kappa) \rightarrow \infty \text{ as } N_\kappa \rightarrow 1.$$

From (2.10), we derive the constraint

$$0 < \kappa < 1. \tag{2.11}$$

We have the following monotonicity properties of N_κ and U_κ w.r.t. κ .

Theorem 2.2 (Monotonicity of U_κ and $E_\kappa^{(0)}$). *If $0 < \kappa_1 < \kappa_2 < 1$, then*

(1) *the monotonicity of U_κ w.r.t. κ holds point-wisely*

$$U_{\kappa_1}(x) > U_{\kappa_2}(x), \quad x \in (0, \pi/2]; \quad (2.12)$$

(2) *the monotonicity of $E_\kappa^{(0)}$ w.r.t. κ also holds*

$$E_{\kappa_1}^{(0)} < E_{\kappa_2}^{(0)}. \quad (2.13)$$

Moreover,

$$\lim_{\kappa \rightarrow 0} \frac{E_\kappa^{(0)}}{\kappa} = \frac{4}{3} \sqrt{2}. \quad (2.14)$$

Proof. The proof of monotonicity can be found in [10]. Here we only show the proof of (2.14). For $0 < \kappa \ll 1$, it can be proved (see [10]) that U_κ satisfies

$$0 \leq \tanh\left(\frac{x}{\sqrt{2\kappa}}\right) - U_\kappa(x) \leq \exp\left(-\frac{c}{\kappa}\right), \quad \forall 0 \leq x \leq \frac{\pi}{2}, \quad (2.15)$$

where $c > 0$ is an absolute constant. Then (2.14) follows from a simple computation

$$\lim_{\kappa \rightarrow 0} \frac{E_\kappa^{(0)}}{\kappa} = \sqrt{2} \int_{\mathbb{R}} (\tanh^2 y - 1)^2 dy = \sqrt{2} \int_{\mathbb{R}} (1 - \tanh^2 y) d(\tanh y) = \frac{4}{3} \sqrt{2}. \quad (2.16)$$

This completes the proof of (2.14). \square

The monotonicity of ground state energy w.r.t. κ , the limit property (2.14), and the monotonicity of U_κ w.r.t. κ , are verified numerically in Fig. 3.

Note that instead of solving the Allen-Cahn dynamics it is possible to compute the profile of the aforementioned ground state U_κ in two straightforward ways, namely, determine N_κ via (2.10) by using Newton's iteration method, or solve the ODE (2.9) defined on $[0, \frac{\pi}{2}]$ by using some standard ODE solvers such as the second-order predictor-corrector method.

3 Symmetry breaking

The following sharp convergence result is proved in [10].

Theorem 3.1 (Convergence for single bump initial data). *Let $0 < \kappa < 1$. Assume the initial data $u_0: [-\pi, \pi] \rightarrow \mathbb{R}$ is odd, 2π -periodic, and $E(u_0) \leq \frac{\pi}{2}$. Suppose u_0 is monotonically increasing on $[0, \frac{\pi}{2}]$ and $u_0(\pi - x) = u_0(x)$ for all $\frac{\pi}{2} \leq x \leq \pi$. Then $u(x, t) \rightarrow U_\kappa$ as $t \rightarrow \infty$.*

In particular it follows from Theorem 3.1 that the solution corresponding to $u_0(x) = \sin(x)$ should converge to a nontrivial odd steady state. However, quite surprisingly, the following numerical computation shows that this is not always the case.




Figure 3: $E_\kappa^{(0)}$ w.r.t. $\kappa \in [0,1]$ and the ground state U_κ for different κ .

3.1 A prototypical example on loss of parity

We solve the following Allen-Cahn dynamics

$$\begin{cases} \partial_t u = \kappa^2 \partial_{xx} u - f(u), \\ u_0 = \sin(x), \end{cases} \quad (3.1)$$

using the first-order implicit-explicit scheme

$$\frac{u^{n+1} - u^n}{\tau} = \kappa^2 \partial_{xx} u^{n+1} - f(u^n), \quad n \geq 0, \quad (3.2)$$

with $\kappa = 0.9$ and time step $\tau = 0.01$. More precisely, we use the pseudo-spectral method (see, e.g., [11, 13] with $N = 256$ Fourier modes for space discretization. The computed steady state and its energy evolution is illustrated in Fig. 4. It is observed that the oddness is lost gradually in time and the computed steady state is completely incorrect.

3.2 Amplification of machine error

We suspect that the above unreasonable symmetry-breaking phenomenon is connected with the machine truncation error. To understand this, one can consider a simple testing ODE

$$u'(t) = u(t), \quad u_0 = 10^{-15} \approx 0, \quad (3.3)$$

where $u_0 \ll 1$ is regarded as the small machine error. Clearly, the solution is $u(t) = u_0 \exp(t)$ so that $u(35) \approx 1.5860 > 1$. However, if $u_0 = 0$ rigorously, $u(t) \equiv 0$. This means that the initial machine error is amplified significantly after $t > 35$.




Figure 4: Wrong steady state (red curves) for (3.1) computed by the first-order IMEX scheme with $\kappa = 0.9$, $\tau = 0.01$, and $N = 256$.




Figure 5: Initial data of $u_0 = \sin(8x)$ at zero points respectively on $[-\pi, \pi]$ and $[0, 2\pi]$, computed by MATLAB. Symmetries at $(0, 0)$ (left) and $(\pi, 0)$ (right) are not preserved due to machine errors.

A further investigation shows that the data stored by computer can be easily “polluted”, even at time $t = 0$. For example, if we input $u_0(x) = \sin(8x)$, we have $u_0(\frac{\pi j}{8}) = 0$ for all integers j . However, standard software such as MATLAB gives nonzero values due to machine errors. As a consequence, the oddness is already violated at time $t = 0$. See Fig. 5 for a graphical illustration. This suggests that the values of u may be contaminated by machine round-off errors at the very beginning of simulation.

Even if we rectify the initial data to be odd (by numerically forcing u_0 to be odd), the oddness can not be ensured after a few iterations, due to the gradual accumulation of




Figure 6: Amplification of machine error of the testing ODE (3.3) (left); an example of the evolution of $u(0,t)$ for the Allen-Cahn equation (1.1) with odd initial condition (right).

machine error. This issue is serious, since it leads to convergence to the incorrect steady state and thereby destroys the fidelity of the numerical simulation in the longer run. In a deeper way the problem of the spurious growth of $u(0,t)$ (see Fig. 6) is connected with the amplification of projections into the exponentially-fast directions (such as projections into the steady state functions ± 1 for the Allen-Cahn case) which can be summarized in the following abstract diagram in Fig. 7.




Figure 7: Schematic diagram of the amplification of machine error.

4 Our filtered approach

4.1 A new symmetry-preserving filter

It is known that $u(x,t)$ should preserve the oddness for all t if the initial condition u_0 is odd. However, standard numerical methods may converge to the spurious steady states in long time due to the accumulation of machine error. To resolve this issue we introduce a new symmetry-preserving filter which amounts to forcing the parity of the solution at every iteration step. Assume the initial data u_0 is odd and has 2π as its minimal period. To preserve the oddness and the invariance of Fourier sine series, we propose a new Fourier filter imposing the zeroth Fourier coefficient to be zero and the real part of all




Figure 8: Correct steady state (red curves) for problem (3.1) computed by the filtered first-order IMEX scheme with $\kappa=0.9$, $\tau=0.01$, and $N=256$.

Fourier coefficients to remain zero, namely, $\forall n \geq 0$,

$$\widehat{u}^n(k=0) = 0; \quad (4.1a)$$

$$\text{Re}(\widehat{u}^n(k)) = 0. \quad (4.1b)$$

One should note that in a typical MATLAB implementation, for (4.1) to hold, we just need to use the command

$$\widehat{u}^n(k) = \text{Im}(\widehat{u}^n(k)) * 1i \quad (4.2)$$

in each iteration. Fig. 8 illustrates the steady state computed with this Fourier filter respectively, where $u_0 = \sin(x)$, $\kappa = 0.9$, $T = 100$. Compared to the result without filter in Fig. 4, the proposed filter clearly helps the algorithm to converge to the correct steady state in long time.

Further, we illustrate plots of $\max_x |u_\infty|$ w.r.t. κ in Fig. 9, computed respectively by the IMEX Fourier scheme (3.2) without and with filter. It can be seen that for the classical method without filter, $\max_x |u_\infty| = 1$ when $\kappa \gtrsim 0.70$, indicating that the computed steady state is wrong. On the other hand, the filtered method guarantees that the numerical solution converges to the correct steady state.

4.2 Convergence of the filtered approach

In this part, we give some mathematical analysis to demonstrate that the filtered numerical solution can converge to the desired steady state solution.

To explain the heart of the matter, we shall carry out the analysis for a semi-discrete model with perturbations introduced by machine error. To simplify the analysis we only consider a restricted set of parameters. For example, we take the representative case




Figure 9: $\max_x |u_\infty|$ w.r.t. κ computed by the first-order IMEX scheme respectively without filter (left) and with filter (right).

$\kappa = 0.9$ which was considered in the earlier numerical experiments (see Figs. 4 and 8) to ensure the uniqueness of the zero-up ground state and simplify the relevant perturbative analysis. To minimize technicality we did not work with the optimal assumptions. In forthcoming works we will develop a comprehensive theoretic framework to address all issues related to full discretization, truncation error and so on.

To this end, we consider $\kappa = 0.9$ and

$$\frac{w-u}{\tau} = \kappa^2 \partial_{xx} w - (u^3 - u), \quad \text{on } [-\pi, \pi], \quad (4.3)$$

where $u, w: \mathbb{T} \rightarrow \mathbb{R}$ are both odd and smooth. Clearly

$$w = (1 - \kappa^2 \tau \partial_{xx})^{-1} [u - \tau(u^3 - u)]. \quad (4.4)$$

We denote the map $u \mapsto w$ as

$$w = \mathcal{N}(u). \quad (4.5)$$

The main model for the filtered solution is given by

$$\begin{cases} u^{n+1} = \mathcal{N}(v^n), \\ v^{n+1} = u^{n+1} + \varepsilon^{n+1}, \\ v^0 = u^0 + \varepsilon^0, \end{cases} \quad (4.6)$$

where $\varepsilon^n: \mathbb{T} \rightarrow \mathbb{R}$ are given machine error functions. Our standing assumption is that

$$\sup_{n \geq 0} \|\varepsilon^n\|_{H^1(\mathbb{T})} \leq \varepsilon_* \ll 1, \quad (4.7)$$

where $\varepsilon_* > 0$ is a very small constant which corresponds to machine precision. We shall assume that all ε^n are odd. This very assumption corresponds to our filtering technique, i.e., we apply the Fourier filter at each iteration step. One should note that u^{n+1} corresponds to the exact numerical solution computed flawlessly (i.e. without machine error) using the numerical scheme, and v^{n+1} corresponds to the actual numerical solution taking into account of the machine error.

We point out that the H^1 assumption (4.7) is reasonable at least for moderately small N , where N is the number of Fourier modes in the Fourier collocation method. A more realistic assumption is to use only L^∞ -norm. Here to simplify the relevant analysis we employ this slightly stronger assumption.

In Lemma 4.1 below, we just consider the case of $\kappa = 0.9$, which ensures that there exists a unique odd zero-up ground state. Moreover, one should note that as $\kappa \rightarrow 1$, the energy of the odd zero-up ground state goes to $\frac{\pi}{2}$ (see Fig. 3) and the constant C_1 in (4.10) has to be taken sufficiently close to zero. In order not to overburden the reader with these technicalities and to simplify the analysis, we consider the simplest case of $\kappa = 0.9$ to illustrate the main ideas.

Lemma 4.1 (Almost steady state). *Fix $\kappa = 0.9$ and assume that U_{steady} is the unique odd zero-up ground state for the equation $\kappa^2 u'' + u - u^3 = 0$ on $\mathbb{T} = [-\pi, \pi]$. Suppose $u: \mathbb{T} \rightarrow \mathbb{R}$ is a smooth odd function satisfying*

$$\kappa^2 u'' + u - u^3 = r(x), \quad (4.8)$$

where the residual error function $r(x)$ satisfies

$$\|r\|_2 \leq r_* \ll 1. \quad (4.9)$$

Assume

$$E(u) = \int_{\mathbb{T}} \left(\frac{1}{2} \kappa^2 (u')^2 + \frac{1}{4} (u^2 - 1)^2 \right) dx \leq \frac{\pi}{2} - C_1, \quad C_1 = 0.001. \quad (4.10)$$

Suppose $u'(0) \geq 0$, then

$$\|u - U_{\text{steady}}\|_{H^1(\mathbb{T})} \leq \alpha_1 r_*, \quad (4.11)$$

where $\alpha_1 = \mathcal{O}(1)$ is an absolute constant.

Proof. We only sketch the details. Denote $u'(0) \geq 0$ and observe that we have $\|u'\|_\infty = \mathcal{O}(1)$. Consider

$$\begin{cases} \kappa^2 u'' + u - u^3 = r, \\ u(0) = 0, \quad u'(0) = p, \end{cases} \quad \begin{cases} \kappa^2 w'' + w - w^3 = 0, \\ w(0) = 0, \quad w'(0) = p. \end{cases} \quad (4.12)$$

Note that here we use the letter p to denote $u'(0)$ which is given (i.e. p is not a parameter). Since u and w have the same initial conditions, we obtain that

$$\|u - w\|_{H^1(\mathbb{T})} \leq \mathcal{O}(1)r_*, \quad |E(u) - E(w)| \ll 1. \quad (4.13)$$

Clearly $E(w) \leq \frac{\pi}{2} - 0.001$. By using the classification of steady state [10], we obtain that w must coincide with U_{steady} . In particular, this shows that $p = u'(0)$ in (4.12) cannot be arbitrary and has to coincide with the ground state. The desired result follows easily. \square

Lemma 4.2. Fix $\kappa = 0.9$ and assume U_{steady} is the unique odd zero-up ground state for the equation

$$\kappa^2 u'' + u - u^3 = 0 \quad \text{on } \mathbb{T} = [-\pi, \pi].$$

Suppose that $u \in H^1(\mathbb{T})$ is an odd function on \mathbb{T} with

$$E(u) = \int_{\mathbb{T}} \left(\frac{1}{2} \kappa^2 (u')^2 + \frac{1}{4} (u^2 - 1)^2 \right) dx \leq \frac{\pi}{2} - C_1, \quad C_1 = 0.001. \quad (4.14)$$

Then we have

$$\min \{ \|u - U_{\text{steady}}\|_{H^1(\mathbb{T})}, \|u + U_{\text{steady}}\|_{H^1(\mathbb{T})} \} \leq C_2 \sqrt{E(u) - E(U_{\text{steady}})}, \quad (4.15)$$

where $C_2 > 0$ is an absolute constant.

Proof. This follows from a general (nontrivial) spectral estimate established in [10]. \square

With the above two lemmas, we now present the main theorem of this section.

Theorem 4.1. Assume the initial state $u^0: \mathbb{T} \rightarrow \mathbb{R}$ is odd, $\|u^0\|_{\infty} \leq 1$ and

$$E(u^0) < \frac{\pi}{2} - 0.002. \quad (4.16)$$

Suppose that the machine error functions $\{\varepsilon^n\}_{n \geq 0}$ are odd and satisfy (4.7). Let $\kappa = 0.9$ and consider (4.6) with $\tau_* \leq \tau \leq \frac{1}{2}$, where $\tau_* = \sqrt{\varepsilon_*}$. Then the following hold.

$$(1) \quad E(U_{\text{steady}}) \leq E(v^{n+1}) \leq \frac{\pi}{2} - 0.001, \quad \forall n \geq 0.$$

(2) For n sufficiently large, we have

$$\min \left\{ \|v^n - U_{\text{steady}}\|_{H^1(\mathbb{T})}, \|v^n + U_{\text{steady}}\|_{H^1(\mathbb{T})} \right\} \leq \tilde{\alpha} \varepsilon_*^{\frac{1}{8}}, \quad (4.17)$$

where $\tilde{\alpha} = \mathcal{O}(1)$ is an absolute constant.

Proof. By a recent result in [9], for any $\tau_* \leq \tau \leq \frac{1}{2}$, we have

$$\sup_{n \geq 0} \max \{ \|u^n\|_\infty, \|v^n\|_\infty \} \leq 1.1.$$

We begin with the basic energy inequality. Namely for any $\tau_* \leq \tau \leq \frac{1}{2}$ and $w = \mathcal{N}(u)$ with u odd, $E(u) \leq \frac{\pi}{2} - 0.001$ and $\max \{ \|w\|_\infty, \|v\|_\infty \} \leq 1.1$, we have

$$\frac{1}{5\tau} \|w - u\|_2^2 + \frac{1}{2} \kappa^2 \|\partial_x(w - u)\|_2^2 + E(w) \leq E(u). \quad (4.18)$$

The main inductive assumption is

$$E(v^n) \leq \frac{\pi}{2} - 0.001. \quad (4.19)$$

Clearly, this holds at the base step $n = 0$. We now assume it holds at step n and consider the solution at $n + 1$ given by

$$v^{n+1} = u^{n+1} + \varepsilon^{n+1} = \mathcal{N}(v^n) + \varepsilon^{n+1}. \quad (4.20)$$

Now we discuss two cases.

Case 1: $\|u^{n+1} - v^n\|_2 / \tau \leq \delta_*^{(1)}$, where $\delta_*^{(1)} \ll 1$ is some suitably small constant to be specified at the end of the proof. Note that by using $u^{n+1} = \mathcal{N}(v^n)$ and the fact that $E(v^n) \leq \frac{\pi}{2} - 0.001$, we have

$$E(u^{n+1}) \leq E(v^n) \quad \text{and} \quad \|u^{n+1}\|_{H^1(\mathbb{T})} = \mathcal{O}(1). \quad (4.21)$$

Thus

$$\kappa^2 \partial_{xx} u^{n+1} - f(u^{n+1}) = \frac{u^{n+1} - v^n}{\tau} + f(v^n) - f(u^{n+1}) = r^{n+1}(x), \quad (4.22)$$

where $\|r^{n+1}\|_2 \ll 1$. By Lemma 4.1 we obtain

$$\min \left\{ \|u^{n+1} - U_{\text{steady}}\|_{H^1(\mathbb{T})}, \|u^{n+1} + U_{\text{steady}}\|_{H^1(\mathbb{T})} \right\} \leq \alpha \delta_*^{(1)}, \quad (4.23)$$

where $\alpha > 0$ is an absolute constant (note that in the more general case, α should depend on κ). Thus in this case we have

$$\min \left\{ \|v^{n+1} - U_{\text{steady}}\|_{H^1(\mathbb{T})}, \|v^{n+1} + U_{\text{steady}}\|_{H^1(\mathbb{T})} \right\} \leq \alpha \delta_*^{(1)} + \varepsilon_*. \quad (4.24)$$

Clearly then $E(v^{n+1}) \leq \frac{\pi}{2} - 0.001$.

Case 2: $\|u^{n+1} - v^n\|_2 / \tau > \delta_*^{(1)}$. By Eq. (4.18) we have

$$\begin{aligned} E(v^n) &\geq E(u^{n+1}) + \frac{1}{5\tau} \|u^{n+1} - v^n\|_2^2 \\ &\geq E(u^{n+1}) + \frac{1}{10\tau} (\delta_*^{(1)})^2 \tau^2 + \frac{1}{10\tau} \|u^{n+1} - v^n\|_2^2 \\ &\geq E(v^{n+1}) + \mathcal{O}(\varepsilon_*) + \frac{1}{10} (\delta_*^{(1)})^2 \tau + \frac{1}{10\tau} \|u^{n+1} - v^n\|_2^2. \end{aligned} \quad (4.25)$$

Thus if

$$\mathcal{O}(\varepsilon_*) + \frac{1}{10} (\delta_*^{(1)})^2 \tau > 0, \quad (4.26)$$

then we obtain

$$E(v^n) \geq E(v^{n+1}) + \frac{1}{10\tau} \|u^{n+1} - v^n\|_2^2. \quad (4.27)$$

Thus for all cases we have proved (4.19). With (4.19) in hand, we now claim that there exists some $n_0 \geq 1$ such that Case 1 holds at $n = n_0$, i.e.

$$\frac{\|u^{n+1} - v^n\|_2}{\tau} \leq \delta_*^{(1)}. \quad (4.28)$$

Assume this is not true, then we are always in Case 2. By (4.27) we obtain

$$\frac{1}{10\tau} \sum_{n=1}^{\infty} \|u^{n+1} - v^n\|_2^2 < \infty.$$

This clearly implies the existence of n_0 . Thus the claim (4.28) holds.

Next, we show that if for some n_0 , (4.28) holds, then the energy of all future v^{n+1} , $n \geq n_0$ will remain in the vicinity of the ground state, i.e.

$$E(v^{n+1}) \leq E(U_{\text{steady}}) + C_3 \cdot (\alpha \delta_*^{(1)} + \varepsilon_*), \quad (4.29)$$

where $C_3 > 0$ is an absolute constant. Indeed, (4.29) holds for $n = n_0$ and any $m > n_0$ which is in Case 1. Now for any $m > n_0$ such that m is in Case 2. We define

$$m_* = \min\{j: j \leq m \text{ and any } \tilde{m} \text{ in } [j, m] \text{ is in Case 2}\}.$$

Clearly by monotonicity of energy in Case 2, we have

$$E(v^{n+1}) \leq E(v^{m_*+1}) \leq E(v^{m_*}). \quad (4.30)$$

Since by definition, $m_* - 1$ is in Case 1, we have

$$E(v^{m_*}) \leq E(U_{\text{steady}}) + C_3 \cdot (\alpha \delta_*^{(1)} + \varepsilon_*), \quad (4.31)$$

thus (4.29) holds for all v^{n+1} , $n \geq n_0$. It follows from Lemma 4.2 that for any $n \geq n_0$,

$$\min \left\{ \|v^{n+1} - U_{\text{steady}}\|_{H^1(\mathbb{T})}, \|v^{n+1} + U_{\text{steady}}\|_{H^1(\mathbb{T})} \right\} \leq C_2 \sqrt{C_3 \cdot (\alpha \delta_*^{(1)} + \varepsilon_*)}. \quad (4.32)$$

To see that (4.32) gives us the desired result, we now specify the parameters used in the proof. We take $\tau_* = \varepsilon_*^{\frac{1}{2}}$ and $\delta_*^{(1)} = \beta \varepsilon_*^{\frac{1}{4}}$ with β being an $\mathcal{O}(1)$ constant, then (4.26) is clearly satisfied. Plugging the values of $\delta_*^{(1)}$ and ε_* into (4.32), we obtain the desired conclusion. \square

We close this section by making several remarks relevant to the above theorem and its proof.

- The condition (4.16) is to ensure that the energy of u^0 is below $\pi/2 = E(0)$ where $E(0)$ is the energy of $u \equiv 0$. This simple assumption avoids the scenario that the solution becomes zero in finite time. For $u^0 = \sin(x)$ and $\kappa = 0.9$, we can compute

$$E(u^0) \approx 1.5327 < \frac{\pi}{2} - 0.002. \quad (4.33)$$

- As we have mentioned earlier, the reasons why we take $\kappa = 0.9$ are: 1) to ensure the uniqueness of the odd zero-up ground state; 2) to simplify the perturbative analysis around the ground state; 3) to back up the numerical experiments (see Figs. 4 and 8). In forthcoming works, we shall treat more general cases.
- To simplify the analysis, we did not optimize the parameters such as τ_* and the other relevant parameters used in the proof. In view of the smallness of the machine error ε_* , the cut-off τ_* is quite a realistic assumption since we usually do not adopt exceedingly small time steps in such long time simulations. Further fine-tuning is certainly possible but we shall not dwell on this issue here.
- From our analysis, it is clear that an improved algorithm can include a stopping criterion when the residual error becomes suitably small. This would also bring drastic simplifications in our proof. In particular, one does not need to consider the situation where the numerical solution hops between Case 1 and Case 2 intermittently.

5 More numerical experiments

In the following numerical experiments, the time step is $\tau = 0.01$ and the number of Fourier modes is $N = 256$ by default. In addition, the “numerical” steady state is obtained when the residual error of the scheme is smaller than $\text{To1} = 10^{-12}$ or the time arrives at $t = 10^5$.

5.1 More 1D Allen-Cahn examples

Example 5.1. We consider more initial conditions. Let the parameter $\kappa = 0.1$ and the initial data $u_0 = \sin(x)$, $\frac{1}{2}(\sin(x) + \sin(2x))$, $\frac{1}{2}(\sin(x) + \sin(4x))$ or $\frac{1}{2}(\sin(x) + \sin(8x))$, respectively.

The approximate steady states are computed using the filtered first-order IMEX pseudo-spectral method. In Fig. 10, it shows that for these odd initial conditions, $u(x,t)$ converges to the same steady state with period 2π .

Example 5.2. We consider more parameter of κ . Let $u_0 = \sin(x)$, $\kappa = 0.999$ and 1.001 respectively.




Figure 10: Example 5.1: Numerical steady states (red curves) for different initial conditions (bold blue curves): (a): $u_0 = \sin(x)$, (b): $\frac{1}{2}(\sin(x) + \sin(2x))$, (c): $\frac{1}{2}(\sin(x) + \sin(4x))$ and (d): $\frac{1}{2}(\sin(x) + \sin(8x))$.




Figure 11: Example 5.2: Computed steady states (red curves) respectively in the cases of (a): $\kappa=0.999$, and (b): $\kappa=1.001$.

It is observed in Fig. 11 that when κ is slightly smaller than 1, the steady state is nonzero, while when κ is slightly larger than 1, the steady state is zero. This is in good agreement with our theory in Section 2.

Example 5.3. We consider an intricate case of metastable states for the Allen-Cahn equation. Let $\kappa = \sqrt{0.001}$ and take two different initial functions $u_0 = \frac{1}{2}(\sin(x) + \sin(2x))$ and $u_0 = \frac{1}{2}(\sin(x) + \sin(8x))$.

It is observed numerically even if the oddness is preserved the numerical solution can get stuck in the metastable states. In particular, it is seen from the left-hand side of Fig. 12 that for $u_0 = \frac{1}{2}(\sin(x) + \sin(2x))$, the corresponding solution u is stuck in the metastable state when the residual error reaches $\text{To1} = 10^{-12}$. Similarly, on the right-hand side for $u_0 = \frac{1}{2}(\sin(x) + \sin(8x))$, the corresponding solution u is stuck in another metastable state. An interesting issue is to investigate these metastable behaviors.

5.2 A 2D Allen-Cahn example

Example 5.4. We consider the 2D Allen-Cahn equation

$$\partial_t u = \kappa^2 \Delta u + u - u^3 \quad \text{on } \mathbb{T}^2 = [-\pi, \pi]^2, \quad (5.1)$$

with $\kappa = 0.1$ and $u_0(x, y) = \sin(x) \sin(y)$.

Clearly, $u(x, y, t)$ should have the form

$$u(x, y, t) = \sum_{k_1, k_2 \geq 1} c_{k_1, k_2}(t) \sin(k_1 x) \sin(k_2 y), \quad (5.2)$$




Figure 12: Example 5.3: Metastable states (red curves) for (a): $u_0 = \frac{1}{2}(\sin(x) + \sin(2x))$ and (b): $u_0 = \frac{1}{2}(\sin(x) + \sin(8x))$, with $\kappa = \sqrt{0.001}$.

where c_{k_1, k_2} is the Fourier coefficient of the $(k_1^{\text{th}}, k_2^{\text{th}})$ mode. This indicates the symmetries $u(x, y, t) = -u(-x, y, t)$ and $u(x, y, t) = -u(x, -y, t)$. In our computation, we use the first-order-IMEX pseudo-spectral method to solve the 2D Allen-Cahn equation with $\tau = 0.01$ and $N_x \times N_y = 256 \times 256$ Fourier modes.

Then, we impose the following 2D symmetry-preserving Fourier filter: $\forall n \geq 0$,

$$\widehat{u}^n(k_1=0, k_2) = 0, \quad \widehat{u}^n(k_1, k_2=0) = 0; \quad (5.3a)$$

$$\text{Im}(\widehat{u}^n(k_1, k_2)) = 0; \quad (5.3b)$$

$$\widehat{u}^n(k_1, k_2) \leftarrow \frac{1}{2}(\widehat{u}^n(k_1, k_2) - \widehat{u}^n(-k_1, k_2)), \quad \forall k_1, k_2; \quad (5.3c)$$

$$\widehat{u}^n(k_1, k_2) \leftarrow \frac{1}{2}(\widehat{u}^n(k_1, k_2) - \widehat{u}^n(k_1, -k_2)), \quad \forall k_1, k_2. \quad (5.3d)$$

In the actual numerical implementation, we apply first (5.3c) and then (5.3d). In MATLAB implementation, we use the following command

```
for j = 1:Ny
un_hat(:, j) = [0; (u_hat(2:end, j) - flip(u_hat(2:end, j)))/2];
end
for i = 1:Nx
u_hat(i, :) = [0, (un_hat(i, 2:end) - flip(un_hat(i, 2:end)))/2];
end
```

It is observed from Fig. 13 that without filter the symmetries are destroyed and u tends to the incorrect steady state. On the other hand, the symmetry can be preserved




Figure 13: Example 5.4: Dynamics of 2D Allen-Cahn equation using the first-order IMEX method without filter (symmetry breaking happens!) where $\kappa=0.1$, $u_0 = \sin(x)\sin(y)$, $\tau=0.01$, $N_x = N_y = 256$.

using this filter, see Fig. 14. A comparison between the corresponding energy evolutions is provided in Fig. 15.

Remark 5.1. It should be pointed out that in this work we single out the odd symmetry as one of the fundamental preserved quantities for two reasons: 1) restriction to odd symmetry eliminates a plethora of unwanted unstable and neutral manifolds; 2) this is the simplest yet unsolved (before our work) scenario where accumulation of machine errors can lead to abrupt escape to spurious energy minimizing states. Our work is of fundamental importance for future investigations in this direction. Even for the 1D simple equation as (1.1), the situation for general initial data remains elusive and one should not take for granted the evenness or oddness as fundamental should-be-conserved symmetries. For example if we take $u_0 = \cos(x) = \sin(x + \frac{\pi}{2})$, then according to our recent classification of steady states in [10], the correct steady state should be $U_\kappa(x + \frac{\pi}{2})$. In this case, the fundamental axis of (odd) symmetry should be $x = -\pi/2$, i.e. the symmetry $u(x) = -u(-\pi - x)$ should be preserved in time. It is possible to develop the corresponding filter and apply our framework to this case, however we shall not dwell on this issue here.




Figure 14: Example 5.4: Same as Fig. 13, except with filter (5.3).

6 Concluding remarks

In this work, we investigated the numerical computation and analysis for steady states of the Allen-Cahn equation with double well potential. Quite surprisingly we find that for very smooth odd initial condition, the numerical solution computed using standard algorithms and very high precision may lose parity in not very long time simulations and eventually converge to the spurious steady state. We call it (numerical) breakdown of symmetry for Allen-Cahn as a manifestation of the gradual accumulation of machine round-off errors. This new phenomenon shows that, just much like other physical conservation laws, conservation of symmetry could possibly be ironed into the fundamental construction of numerical algorithms since the violation of symmetry could lead to completely erroneous long time simulations. To resolve this issue we introduced a new symmetry-preserving filter technique in order to give an accurate long time computation of the steady states. We developed a new theoretical framework taking into account of perturbations introduced by machine errors. This opens doors to many future directions and developments.

In future works, we shall develop a new asymptotic theory and spectral analysis framework around the steady states. We also plan to address the following important issues.




Figure 15: Example 5.4: Energy evolutions of 2D Allen-Cahn equation computed by the first-order IMEX method respectively without filter and with filter.

- General filtering/stabilizing technique for long time numerical computations. In our work we introduced a symmetry-preserving filter to deal with special data with odd symmetry or certain spectral band gaps. In the future we will develop a more systematic filtering procedure to suppress the spurious unstable modes in the long time simulations. In addition, we plan to develop a complete theoretical framework to show the stability of filtered solutions under perturbations by machine errors. This will include full numerical discretization as well as very realistic assumptions on the machine errors and so on.
- Other equations and phase field models. These include nonlocal Allen-Cahn equations driven by general polynomials or logarithmic nonlinearities, and also the Cahn-Hilliard equations, molecular beam epitaxy equation and so on.

Acknowledgments

The research of T. Tang is partially supported by NSFC Grant 11731006, the NSFC/RGC Grant 11961160718. The research of D. Li is supported in part by Hong Kong RGC grant GRF 16307317 and 16309518. The research of W. Yang is supported by NSFC Grants 11801550 and 11871470. The research of C. Quan is supported by NSFC Grant 11901281, the Guangdong Basic and Applied Basic Research Foundation (2020A1515010336), and the Stable Support Plan Program of Shenzhen Natural Science Fund (Program Contract No. 20200925160747003).

References

- [1] G. Alberti, L. Ambrosio, and X. Cabré. On a long-standing conjecture of E. De Giorgi: Symmetry in 3D for general nonlinearities and a local minimality property. *Acta Applicandae Mathematica*, 65(1):9–33, 2001.
- [2] L. Ambrosio and X. Cabré. Entire solutions of semilinear elliptic equations in R^3 and a conjecture of De Giorgi. *Journal of the American Mathematical Society*, 13(4):725–739, 2000.
- [3] M.T. Barlow, R.F. Bass, and C.F. Gui. The Liouville property and a conjecture of De Giorgi. *Communications on Pure and Applied Mathematics*, 53(8):1007–1038, 2000.
- [4] P. Bates, P. Fife, X.F. Ren, and X.F. Wang. Traveling waves in a convolution model for phase transitions. *Archive for Rational Mechanics and Analysis*, 138(2):105–136, 1997.
- [5] H. Berestycki, F. Hamel, and R. Monneau. One-dimensional symmetry of bounded entire solutions of some elliptic equations. *Duke Mathematical Journal*, 103(3):375–396, 2000.
- [6] Y. Cheng, A. Kurganov, Z. Qu, and T. Tang. Fast and stable explicit operator splitting methods for phase-field models. *Journal of Computational Physics*, 303:45–65, 2015.
- [7] M. del Pino, M. Kowalczyk, and J.C. Wei. On De Giorgi’s conjecture in dimension $n \geq 9$. *Annals of Mathematics*, pages 1485–1569, 2011.
- [8] B. Li, J. Yang, and Z. Zhou. Arbitrarily high-order exponential cut-off methods for preserving maximum principle of parabolic equations. *SIAM Journal on Scientific Computing*, 42(6):A3957–A3978, 2020.
- [9] D. Li. Effective maximum principles for spectral methods. *Ann. Appl. Math.*, 37:131–290, 2021.
- [10] D. Li, C. Quan, T. Tang, and W. Yang. Sharp convergence to steady states of Allen–Cahn. *arXiv preprint arXiv:2106.08534*, 2021.
- [11] J. Shen, T. Tang, and L.L. Wang. *Spectral Methods: Algorithms, Analysis and Applications*, volume 41. Springer Science & Business Media, 2011.
- [12] J. Shen, J. Xu, and J. Yang. The scalar auxiliary variable (SAV) approach for gradient flows. *Journal of Computational Physics*, 353:407–416, 2018.
- [13] L.N. Trefethen. *Spectral Methods in MATLAB*. SIAM, 2000.
- [14] K.L. Wang. A new proof of Savin’s theorem on Allen–Cahn equations. *Journal of the European Mathematical Society*, 19(10):2997–3051, 2017.
- [15] C.J. Xu and T. Tang. Stability analysis of large time-stepping methods for epitaxial growth models. *SIAM Journal on Numerical Analysis*, 44(4):1759–1779, 2006.
- [16] J. Xu, Y. Li, S.N. Wu, and A. Bousquet. On the stability and accuracy of partially and fully implicit schemes for phase field modeling. *Computer Methods in Applied Mechanics and Engineering*, 345:826–853, 2019.